Association between a *MYH9* polymorphism (rs3752462) and renal function in the Spanish RENASTUR cohort.

Beatriz Tavira^a, Eliecer Coto^{a,f}, Juan Gómez^a, Salvador Tranche^b, Kevin Miguélez^b,
Francisco Ortega^{c,f}, Beatriz Díez^d, Emilio Sánchez^c, Rafael Marín^c, Jorge Arenas^e,
Victoria Alvarez^a

^a Genética Molecular-Laboratorio Medicina, HUCA, Oviedo, Spain.

^b Centros de Salud Principado de Asturias, Asturias, Spain.

^c Nefrología, HUCA, Oviedo, Spain.

^d Nefrología, Hospital V Nalón, Langreo, Spain

^e CAIBER, HUCA, Oviedo, Spain

^f Fundación Renal and RED INVESTIGACION RENAL (REDINREN), Spain.

Correspondence:

Eliecer Coto García

Genética Molecular

Hospital Universitario Central de Asturias

33006 – Oviedo – Spain

Tel/Fax (34)985107968

Email: eliecer.coto@sespa.princast.es

Keywords: MYH9 polymorphisms, glomerular filtration, renal function

Abstract

The *MYH9* gene encodes a protein that is expressed in the kidney glomerular podocytes. *MYH9* single nucleotide polymorphisms (SNPs) have been linked to the risk for chronic kidney disease (CKD) and end stage renal disease. Our aim was to determine whether *MYH9* SNPs were associated with renal disease in Spanish Caucasians. The RENASTUR cohort consisted of 592 Spanish Caucasians, aged 55-85 years. They were genotyped for SNPs rs3752462 and rs4821480, that tagged haplotype E. The main values between individuals with a glomerular filtration rate (eGFR) <60 and ≥ 60 ml/ min./1.73 m² were statistically compared. The next variables were significantly associated with the eGFR in the univariate analysis: age, gender, type 2 diabetes, total cholesterol, total LDL-cholesterol, and the *MYH9* rs3752462 (TC+TT genotypes; $p=0.003$). This SNP remained significantly associated with the eGFR in the multivariate analysis.

In conclusion, SNP rs3752462 was an independent predictor of reduced eGFR in the Spanish RENASTUR population. The genotyping of this *MYH9* SNP could help to identify individuals at risk of developing CKD.

1. Introduction

The estimated Glomerular filtration rate (eGFR) is commonly used to define renal function and identify individuals with chronic kidney disease (CKD). The Kidney Disease Outcome Quality Initiative (K / DOQI) defined CKD as the presence of renal impairment with an eGFR <60 [National Kidney Foundation, 2002]. Based on this definition the EPIRCE study (Epidemiology of Chronic Kidney Disease in Spain) estimated that approximately 10% of the Spanish adult population had some degree of CKD, and similar values were also estimated from other epidemiological studies [Coresh et al., 2007; Otero et al., 2010]. CKD is a strong predictor of end stage renal disease (ESRD), cardiovascular morbidity, and mortality. In addition to well known risk factors such as hypertension and diabetes, recent genome wide association studies (GWAs) linked several single nucleotide polymorphisms (SNPs) to eGFR and prevalent CKD [Köttgen et al., 2009, 2010]. Some of these SNPs were also associated with the progression to ESRD [Böger et al., 2011].

The *MYH9* gene encodes a myosin heavy chain protein expressed in glomerular podocytes and mesangial cells [Marini et al., 2006]. Recent studies linked *MYH9* SNPs / haplotypes to the risk of developing focal segmental glomerulosclerosis (FSGS), hypertensive nephropathy, and non-diabetic ESRD among African and Hispanic Americans [Kopp et al., 2008; Kao et al., 2008; Freedman et al., 2009a, 2009b; Behar et al., 2010]. This association was not replicated among American Indians, a fact that could be partly explained by differences in the prevalence of risk factors for CKD between the populations [Franceschini et al., 2010].

Among individuals of European ascent, *MYH9* polymorphisms have been linked to kidney function, non-diabetic and diabetic nephropathy, and idiopathic FSGS [Kao et al., 2008; Pattaro et al., 2009; O'Seaghdha et al., 2011; Cooke et al., 2012]. Our objective was to define the effect of common *MYH9* SNPs on renal function in a Spanish cohort of healthy elderly individuals.

2. Material and methods

2.1 Study population and data collection

This study was approved by the Ethical Committee of Hospital Universitario Central de Asturias (HUCA) and all the participants signed an informed consent. The RENASTUR was a cross-sectional study to evaluate the association between gene polymorphisms and renal function in elderly people. A total of 592 individuals aged 55-85, all Caucasian, non-related, and from the region of Asturias (Northern Spain, total population one million), were randomly chosen from the general population in the period 2010-2012. All the participants were evaluated by a qualified physician, who was also responsible for obtaining the main analytical and clinical values (**table 1**).

Smoking (ever/never smoked) was self reported. Body mass index (BMI) was defined by weight and height measured at the exam. Blood pressure values were obtained at exam. Blood glucose, serum creatinine (SCr), total serum cholesterol, high and low density lipoprotein cholesterol, and triglycerides were measured from blood samples collected by venipuncture. Individuals with a documented history of hypertension or diabetes or who were treated with antihypertensive or antidiabetic drugs were classified as hypertensives and diabetics, respectively. The biochemical profiles from fasting blood samples were obtained by routine methods. The estimated glomerular filtration rate (eGFR) was calculated with the Modification of Diet in Renal Disease (MDRD) formulae [Levey et al., 1999].

2.2. MYH9 genotyping

DNA was obtained from 5 ml of blood leukocytes, and two *MYH9* SNPs (rs3752462 and rs4821480) were genotyped through a Polymerase Chain Reaction-restriction fragment length polymorphism (PCR-RFLP) method (**supplementary figure 1**). These SNPs were chosen because they defined four haplotypes (E1-E4) that have been linked to nephropathies [Kao et al., 2008; Kopp et al., 2008; Oleksyk et al., 2010]. Briefly, for the rs3752462 the DNA a 421 bp fragment was amplified with primers

CCAGGAGCATCCGGGCTCTA (forward) and CACCTCCACAACCAACACAGAGCT (reverse). After digestion with *RsaI*, reactions were electrophoresed on 3% agarose gels a constant band and the two alleles were visualized as bands of 243+178 bp (C allele) and 243+92+86 bp (T allele). For the rs4821480 the DNA was PCR amplified with primers CCGCTGGGCAGGGGTGTT (forward) and TCTTCTGTGAGGTTGGT GGTG (reverse), followed by digestion with *DraI* and electrophoresis on agarose gels to visualize the two alleles of 537 bp (G allele) and 322+215 bp (T allele). To confirm the accuracy of this genotyping method, PCR fragments from several individuals with each of the RFLP-genotypes were sequenced using BigDye chemistry in an ABI3130 automated system (Applied Biosystems).

2.3. Statistical analysis

CKD cases were defined as those with an eGFR < 60 ml/min/1.73 m², while healthy controls were those free of CKD at baseline. All the statistical analyses were done with the SPSS statistical software (version 17.0; SPSS Inc., Chicago, IL). The Fisher's exact and student's t tests were used to compare categorical and continuous variables between the groups, respectively. The multivariate regression analysis was performed to determine the independent effect of the variables. P values <0.05 were considered statistically significant. Haplotype frequencies were calculated online (cubic exact solutions for the estimation of pairwise haplotype frequencies; <http://www.oege.org/software/cubex>).

3. Results

Table 1 summarizes the main characteristics in the whole RENASTUR cohort, and in those with $\text{eGFR} < 60$ ($n=77$) and ≥ 60 ($n=515$) ml/min/1.73 m^2 . The genotype frequencies for the two *MYH9* SNPs did not deviate from Hardy–Weinberg equilibrium in the two groups.

In the univariate analysis the next variables were significantly associated with an $\text{eGFR} < 60$: age ($p < 0.001$), male ($p = 0.04$), type 2 diabetes ($p < 0.001$), total cholesterol ($p = 0.03$), LDL-cholesterol ($p = 0.05$), and *MYH9* rs3752462 T allele (dominant effect, TC+TT genotypes; $p = 0.003$). Hypertensives and smokers were non-significantly more frequent in the $\text{eGFR} < 60$ group. In the multivariate logistic regression analysis age, diabetes, gender, and the *MYH9* rs3752462 genotype remained as independent risk factors for $\text{eGFR} < 60$ ml/min/1.73 m^2 (**table 2**). *MYH9* rs3752462 T-carriers had lower mean eGFR compared to CC homozygotes (77 ± 18 vs. 83 ± 19 ; $p = 0.02$). The *MYH9* rs4821480 G allele was more frequent in the $\text{eGFR} < 60$ group (0.12 vs. 0.08), although the difference was not statistically significant. The most common E2 haplotype (rs4821480 T / rs3752462 C) showed a significantly higher frequency in the $\text{eGFR} \geq 60$ group ($p = 0.04$) (**table 3**). This was in agreement with the reported protective effect for this haplotype among Caucasians.

To determine whether the rs3752462 SNP had an effect on pre-mRNA splicing we amplified the cDNA from six heart tissues with primers that matched exons 12 and 14. Only a single PCR band was visualized from all the tissues, included the two that were heterozygotes for this SNP. Sequencing of the amplified fragments showed a normal transcript in all the samples (**supplementary figure 2**). This suggested that the association between rs3752462 and CKD was likely not due to a direct effect pre-mRNA splicing.

4. Discussion

CKD is a major cause of morbidity and mortality, and is thus considered an important community health problem. According to the EPIRCE study approximately 10% of the Spanish adult population had some degree of CKD [Otero et al., 2010]. This was in agreement with our findings in the RENASTUR cohort, with 10% of the participants showing an $\text{eGFR} < 60 \text{ ml/min/1.73 m}^2$. The search for markers associated with CKD is important to identify individuals at risk for ESRD and cardiovascular events, that could benefit from preventive approaches. It is a well established fact that renal function declines with age [Fox et al., 2004]. In addition, several gene polymorphisms have been linked to impaired renal function and CKD. Mutations in the *MYH9* gene cause the Fechtner and Epstein syndromes, both resulting in nephritis to varying degrees [Seri et al., 2000; Arrondel et al., 2002; Bostrom et al., 2010].

Recent studies linked *MYH9* SNPs / haplotypes to the risk of developing FSGS, hypertensive nephropathy, diabetic and non-diabetic nephropathy, and lupus nephritis [Kopp et al., 2008; Kao et al., 2008; Freedman et al., 2009a, 2009b; Behar et al., 2010; Kao et al., 2008; Pattaro et al., 2009; O'Seaghdha et al., 2011; Cooke et al., 2012; Lin et al., 2012]. We found a significant association between rs3752462 and a reduced eGFR among individuals without a previous diagnosis of CKD. In the multivariate analysis this association was independent of other risk factors, such as age and diabetes. The *MYH9* SNP rs4821480 was previously linked to CKD among individuals of African ancestry but was not associated with a reduced eGFR in our cohort. Due to the very low frequency (0.08) of the risk allele in our population, to exclude an association to CKD a much large number of individuals with impaired renal function should be required: for a power of 80 at a $p=0.05$, a total of 4,213 individuals (548 with an $\text{eGFR} < 60$). Haplotype frequencies were similar to the reported from other Caucasian populations [Oleksyk et al., 2010]. In agreement with previous findings the most common E2 haplotype showed a protective effect in our population.

The association between rs3752462 and CKD has been replicated in several Caucasian populations, but a study based on American Indians did not find a significant effect of *MYH9* SNPs/haplotypes on CKD. Because *MYH9* genetic susceptibility has been described mostly for non-diabetic CKD, it is possible that this gene is not an important genetic risk factor for CKD in populations with a high prevalence of T2DM, such as American Indians. Differences in linkage disequilibrium between rs3752462 and the actual functional genetic variation responsible for the associations could also explain the discrepancies between the studies (Franceschini et al., 2010).

Recently, two SNPs in the *APOLI* gene (rs73885319 and rs60910145, haplotype G1) and one deletion (rs71785323, haplotype G2) were linked to CKD/ESRD among individuals of African ascent [Genovese et al., 2010]. These were missense amino-acid changes that could alter the function of ApoL1. Although these *APOLI* polymorphisms could explain part of the effect previously attributed to *MYH9* among individuals of African ascent, the risk alleles were absent among Caucasians [Cooke et al., 2012]. Asturias is a region in the Northern Spain Cantabrian Cornice, and mitochondrial haplogroup-based studies showed a typical European genetic background with no evidence for recent African colonization [García et al., 2010; Palacin et al., 2011]. We thus considered very unlikely that African *APOLI* variants were responsible for the observed effect of *MYH9* SNPs on CKD. To determine whether some *APOLI* variants were linked to CKD in our population this gene should be sequenced in individuals with the *MYH9*-risk genotypes.

SNP rs3752462 was in intron 13 of *MYH9*, and a direct effect on pre-mRNA splicing has not been reported. Amplification of transcripts from heart tissues heterozygotes for this SNP with primers that matched exons 12 and 14 showed a single PCR band that corresponded to the normal *MYH9* sequence. The fact that no aberrant transcripts were amplified suggested a lack of effect on pre-mRNA splicing. The association between this polymorphism and CKD could be explained by linkage disequilibrium

with a functional variant in *MYH9* or a nearby gene. Nelson et al. genotyped 79 *MYH9* SNPs in a total of 2,496 cases (FSGS, HIV-associated nephropathy, and ESRD attributed to hypertension) and controls, and mapped the strongest kidney disease association to the region of introns 13-15 [Nelson et al., 2010]. Neither the highly associated SNPs nor the results of resequencing *MYH9* revealed non-synonymous changes that could account for the disease associations, although some of the SNPs were predicted to affect pre-mRNA splicing and could thus have a functional effect on gene expression and protein function [Nelson et al., 2010].

Finally, although the rs3752462 T allele (CT+TT genotypes) and $\text{eGFR} < 60 \text{ ml/min/1.73 m}^2$ was statistically significant and the sample sizes were enough to reach a power of 80, our work was based on a limited number of individuals (and in particular for the group with an $\text{eGFR} < 60$) and from a single Caucasian population. The replication of this association on larger cohorts from other populations are necessary to confirm or refute this *MYH9* SNP as an independent predictor of renal function.

In conclusion, SNP rs3752462 was related with a decreased eGFR in the RENASTUR population. Because CKD is often silent until advanced stages of disease, many individuals are unaware they suffer from renal disease until symptomatic uremia is detected. Our study suggested that *MYH9* genotyping could help to identify individuals at risk of developing CKD.

Financial disclosure. All the authors declare they have no conflicts of interest related with this work.

Contributorship. All the authors contributed to this work by recruiting the population cohort (S.T., K.M., M.I.R., F.O., B.D., R.M., E.S., J.A.) or performing the genetic and statistical analysis (B.T., J.G., E.C., V.A).

Acknowledgements. This work was supported by grant Red de Investigación Renal-REDINREN, and Fundación Investigación Científica y Técnica (FICYT)-European FEDER funds. This work was partly supported by grant FIS PI10/01971 (F.O.).

References

- Arrondel** C, Vodovar N, Knebelmann B, et al. Expression of the nonmuscle myosin heavy chain IIA in the human kidney and screening for MYH9 mutations in Epstein and Fechtner syndromes. *J Am Soc Nephrol* 2002;13:65-74.
- Behar** DM, Rosset S, Tzur S et al. African ancestry allelic variation at the MYH9 gene contributes to increased susceptibility to non-diabetic end-stage kidney disease in Hispanic Americans. *Hum Mol Genet* 19: 1816–1827.
- Böger** CA, Gorski M, Li M et al. Association of eGFR-related loci identified by GWAS with incident CKD and ESRD. *PLoS Genet* 2011;7:e1002292.
- Bostrom** MA, Freedman BI. The spectrum of MYH9-associated nephropathy. *Clin J Am Soc Nephrol* 2010;5:1107-13.
- Cooke** JN, Bostrom MA, Hicks PJ et al. Polymorphisms in MYH9 are associated with diabetic nephropathy in European Americans. *Nephrol Dial Transplant* 2012;27:1505-11.
- Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, Van Lente F, Levey AS. Prevalence of chronic kidney disease in the United States. *JAMA*. 2007; 298: 2038-47.
- Fox** CS, Larson MG, Leip EP, et al. Predictors of new- onset kidney disease in a community based population. *JAMA* 2004;291:844-50.
- Franceschini** N, Voruganti VS, Haack K, et al. The association of the **MYH9 gene** and kidney outcomes in American Indians: the Strong Heart Family Study. *Hum Genet*. 2010 Mar;127(3):295-301.
- Freedman** BI, Hicks PJ, Bostrom MA, et al. Non-muscle myosin heavy chain 9 gene MYH9 associations in African Americans with clinically diagnosed type 2 diabetes mellitus-associated ESRD. *Nephrol Dial Transplant*. 2009a; 24: 3366-71.

Freedman BI, Kopp JB, Winkler CA et al. Polymorphisms in the nonmuscle myosin heavy chain 9 gene (MYH9) are associated with albuminuria in hypertensive African Americans: the HyperGEN study. *Am J Nephrol* 2009b; 29: 626–32.

García O, Fregel R, Larruga JM, et al. Using mitochondrial DNA to test the hypothesis of a European post-glacial human recolonization from the Franco-Cantabrian refuge. *Heredity* 2011; 106: 37-45.

Genovese G, Friedman DJ, Ross MD, et al. Association of trypanolytic ApoL1 variants with kidney disease in African Americans. *Science* 2010;329: 841–5.

Kao WH, Klag MJ, Meoni LA et al. MYH9 is associated with nondiabetic end-stage renal disease in African Americans. *Nat Genet.* 2008; 40: 1185–92.

Kopp JB, Smith MW, Nelson GW et al. MYH9 is a major-effect risk gene for focal segmental glomerulosclerosis. *Nat Genet* 2008;40:1175-84.

Köttgen A, Glazer NL, Dehghan A et al. Multiple novel loci are associated with indices of renal function and chronic kidney disease. *Nat Genet* 2009; 41: 712–7.

Köttgen A, Pattaro C, Böger CA et al. Multiple New loci associated with kidney function and chronic kidney disease. *Nat Genet* 2010;42:376–84.

Levey AS, Bosch JP, Lewis JB, et al. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 1999;130:461-70.

Lin CP, Adrianto I, Lessard CJ, Kelly JA et al. Role of **MYH9** and APOL1 in African and non-African populations with lupus nephritis. **Genes Immun** 2012 13: 232-8.

Marini M, Bruschi M, Pecci A et al. Non-muscle myosin heavy chain IIA and IIB interact and co-localize in living cells: relevance for MYH9-related disease. *Int J Mol Med* 2006;17:729-36.

National Kidney Foundation. K/ DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002; 39 Suppl 1:46-75.

Nelson GW, Freedman BI, Bowden DW et al. Dense mapping of MYH9 localizes the strongest kidney disease associations to the region of introns 13 to 15. *Hum Mol Genet.* 2010; 19:1805-1815.

Oleksyk TK, Nelson GW, An P, Kopp JB, Winkler CA. Worldwide distribution of the MYH9 kidney disease susceptibility alleles and haplotypes: evidence of historical selection in Africa. *PLoS One.* 2010 5: e11474.

O'Seaghdha CM, Parekh RS, Hwang SJ et al. The MYH9/APOL1 region and chronic kidney disease in European-Americans. *Hum Mol Genet* 2011; 20: 2450–56.

Otero A, De Francisco AL, Gayoso P, et al. Prevalencia de la insuficiencia renal crónica en España: resultados del estudio EPIRCE. *Nefrología* 2010; 30: 78-86.

Palacín M, Alvarez V, Martín M, et al. Mitochondrial DNA and TFAM gene variation in early-onset myocardial infarction: evidence for an association to haplogroup H. *Mitochondrion.* 2011; 11: 176-81.

Pattaro C, Aulchenko YS, Isaacs A et al. Genome-wide linkage analysis of serum creatinine in three isolated European populations. *Kidney Int* 2009; 76: 297–306.

Seri M, Cusano R, Gangarossa S, et al. Mutations in MYH9 result in the May-Hegglin anomaly, and Fechtner and Sebastian syndromes. The May-Hegglin/Fechtner Syndrome Consortium. *Nat Genet* 2000;26:103-5.

Table 1. Main values in the total population, and in individuals with eGFR <60 and ≥ 60 ml/min per 1.73 m^2

Main characteristics	TOTAL Median (range) (n=592)	eGFR <60 [#] Median (range) (n=77)	eGFR >60 Median (range) (n=515)	P values
Age (years)	73(51-83)	76(56-81)	72(51-81)	<0.001
Weight (kg)	75(43- 166)	78(54-153)	74(43-163)	0.10
BMI (kg/m ²)	29.35 (17.63-47.78)	29.72 (22.04-47.78)	29.24 (17.63-46.87)	0.62
Glucose levels (mg/dl)	96(7-324)	97(75-206)	96(7-324)	0.05
Cholesterol (mg/dl)	217(18-335)	205(18-321)	219(71-335)	0.03
LDL (mg/dl)	132(10-249)	125(48-239)	133(41-249)	0.05
HDL (mg/dl)	58(5-276)	53(5-276)	58(25-158)	0.01
Triglycerides (mg/dl)	102(8-470)	119(8-436)	101(39-470)	0.07
eGFR (ml/min per 1.73 m^2)	82 (43-169)			
Creatinine serum levels (mg/dl)	0.80(0.50-76)	1.20(0.70-76)	0.80(0.40-1.30)	<0.001
Albuminuria ratio (mg/g) (albumin/creatinine ratio)	1.6 (0.30-1523)	4.80(0-1522.90)	1.50(0-561.10)	0.001
Male	260 (44%)	42 (55%)	218 (42%)	0.04
Hypertensives	153 (26%)	25 (32%)	128 (25%)	0.15
Smokers	69 (12%)	12 (16%)	57 (11%)	0.25
Diabetics (type II)	169 (29%)	35 (45%)	134 (26%)	<0.001
Hypercholesterolemics	311 (52%)	44 (57%)	267 (52%)	0.39
MYH9 rs3752462				
CC	277/592 (0.47)	25/77 (0.32)	252/515 (0.49)	0.003*
CT	260/592 (0.44)	44/77 (0.57)	216/515 (0.42)	
TT	55/592 (0.09)	8/77 (0.11)	47/515 (0.09)	
C	814 (0.69)	94 (0.61)	720 (0.70)	
T	370 (0.31)	60 (0.39)	310 (0.30)	
MYH9 rs4821480				
TT	497/592 (0.84)	61/77(0.79)	435/515 (0.84)	0.16
TG	95/592 (0.16)	15/77 (0.19)	80/515 (0.16)	
GG	1/592 (<0.01)	1/77 (0.02)	0/515	
T	1,089 (0.92)	137 (0.88)	950 (0.92)	0.17
G	97 (0.08)	17 (0.12)	80 (0.08)	

[#] eGFR (ml/ min./ 1.73 m^2) = $186 \times [\text{plasma creatinin (mg/dl)}] - 1.154 \times (\text{age}) - 0.203 \times (0.742 \text{ if female}) \times (1.212 \text{ if black})$.

* GA + AA vs. GG, Fisher's exac test.

Table 2. Univariate and multivariate binary logistic analysis regression. The dependent variable is eGFR (<60 vs. ≥60 ml/min per 1.73 m²).

	Univariate Analysis P values/ OR (min-max)	Multivariate Analysis P values/ OR (min-max)
Age (years)	<0.001/0.90(0.86-0.95)	<0.001/0.87(0.82-0.92)
Weight (kg)	0.123/0.98(0.97-1)	
BMI (kg/m ²)	0.676/0.98(0.93-1.04)	
Total cholesterol (mg/dl)	0.003/1.009(1.003-1.016)	
LDL (mg/dl)	0.037/1.008(1-1.016)	
HDL (mg/dl)	0.309/1.008(0.99-1.02)	
Triglycerides (mg/dl)	0.09/0.99(0.993-1)	
Male	0.04/0.59(0.36-0.98)	0.02/0.50(0.28-0.90)
HTA (yes)	0.15/1.22(0.70-2.14)	
Smoker (yes)	0.25/1.75(0.99-3.09)	
Diabetes type II (yes)	<0.001/2.26(1.33-3.83)	0.004/2.39(1.32-4.43)
Hypercholesterolemia (yes)	0.39/1.17(0.69-1.99)	
MYH9 rs3752462 *	0.007/1.99(1.20-3.31)	0.046/1.86(1.01-3.42)
MYH9 rs4821480	0.893/1.05(0.47-2.33)	

* GA+AA vs. GG, OR=1.99 (1.20-3.31)

Table 3. Haplotype frequencies (rs4821480 / rs3752462) in our population and in the Vasque and French populations [Oleksyk, 2010].

HAPLOTYPE	TOTAL	eGFR <60	eGFR ≥60	VASQUE	FRENCH
E1 (GT)	0.06	0.05	0.06	0.05	0.05
E2 (TC)	0.65	0.58	0.66	0.69	0.66
E3 (TT)	0.27	0.34	0.26	0.23	0.29
E4 (GC)	0.02	0.02	0.02	-	-

E2: eGFR>60 vs eGFR<60, p=0.04.

MYH9-rs4821480

MYH9-pol3752462

Supplementary Figure 1. PCR-RFLP genotyping of the two *MYH9* polymorphisms (reverse strand nucleotides).

Supplementary figure 2.

Left: Amplification of cDNAs synthesised from *MYH9*-mRNA of heart tissues with primers that matched exons 23 and 24, and 12 and 14 (left).

Right: sequencing of the cDNA amplifications showing the correct exon12-exon13 and exon24-exon25 boundaries in an individual heterozygous for rs3752462.