

A genetic linkage map of *Phaseolus vulgaris* L. and localization of genes for specific resistance to six races of anthracnose (*Colletotrichum lindemuthianum*)

Cristina Rodríguez-Suárez · Belén Méndez-Vigo ·
Astrid Pañeda · Juan José Ferreira · Ramón Giraldez

Received: 27 July 2006 / Accepted: 17 November 2006 / Published online: 22 December 2006
© Springer-Verlag 2006

Abstract A genetic map of common bean was constructed using 197 markers including 152 RAPDs, 32 RFLPs, 12 SCARs, and 1 morphological marker. The map was established by using a F₂ population of 85 individuals from the cross between a line derived from the Spanish landrace Andecha (Andean origin) and the Mesoamerican genotype A252. The resulting map covers about 1,401.9 cM, with an average marker distance of 7.1 cM and includes molecular markers linked to disease resistance genes for anthracnose, bean common mosaic virus, bean golden yellow mosaic virus, common bacterial blight, and rust. Resistance to races 6, 31, 38, 39, 65, and 357 of the pathogenic fungus *Colletotrichum lindemuthianum* (anthracnose) was evaluated in F₃ families derived from the corresponding F₂ individuals. The intermediate resistance to race 65 proceeding from Andecha can be explained by a single dominant gene located on linkage group B1, corresponding to the *Co-1* gene. The recombination between the resistance specificities proceeding from A252 agrees with the assumption that total resistance to races 6, 31, 38, 39, 65, and 357, is organized in two

clusters. One cluster, located on B4 linkage group, includes individual genes for specific resistance to races 6, 38, 39, and 357. The second cluster is located on linkage group B11 and includes individual genes for specific resistance to races 6, 31, 38, 39, and 65. These two clusters correspond to genes *Co-3/Co-9* and *Co-2*, respectively. It is concluded that most anthracnose resistance *Co-* genes, previously described as single major genes conferring resistance to several races, could be organized as clusters of different genes conferring race-specific resistance.

Introduction

Common bean (*Phaseolus vulgaris* L.) is the most important food legume for direct human consumption worldwide (Schoonhoven and Voysest 1991). It is a diploid legume ($2n = 22$) with a relatively small genome (0.65 pg per haploid genome; Aramuganathan and Earle 1991). Several genetic maps have been developed in common bean (Vallejos et al. 1992; Nodari et al. 1993; Adam-Blondon et al. 1994) including different economic traits of interest, such as domestication syndrome, crop quality, tolerance to abiotic stresses, and disease resistances. An integrated linkage map was established in order to align these previous maps (Freyre et al. 1998), new microsatellite markers have been described and located on the integrated linkage map (Blair et al. 2003), and the correspondence between linkage groups (B1–B11) and chromosomes has been established (Pedrosa et al. 2003).

Anthracnose, caused by the hemibiotrophic fungus *Colletotrichum lindemuthianum*, is the most serious disease of common bean worldwide due to its seed-borne

Communicated by F. Muehlbauer.

C. Rodríguez-Suárez and B. Méndez-Vigo equally share for authorship.

C. Rodríguez-Suárez · B. Méndez-Vigo · A. Pañeda ·
R. Giraldez (✉)
Department of Biología Funcional,
University of Oviedo, 33006 Oviedo, Spain
e-mail: giralddez@uniovi.es

J. J. Ferreira
Area de Cultivos Hortofrutícolas, SERIDA,
33300 Villaviciosa (Asturias), Spain

nature and pathogenic variability (Pastor-Corrales and Tu 1994). Currently, races of *C. lindemuthianum* are classified by inoculating isolates onto a universal set of 12 differential cultivars and named based on a binary nomenclature system (Pastor-Corrales 1991). Genetic resistance is regarded as the most effective disease management strategy for the control of anthracnose. Several genes conferring dominant resistance to different sets of pathogenic races have been described and located to the integrated linkage map, either by direct mapping or indirectly, using linked molecular markers: *Co-1* on linkage group B1 (Barrus 1915; Méndez de Vigo 2001), *Co-2* on B11 (Mastenbroek 1960; Adam-Blondon et al. 1994), *Co-3* on B4 (Bannerot 1965; Méndez-Vigo et al. 2005), *Co-4* on B8 (Fouilloux 1976; Melotto et al. 2004; Méndez de Vigo 2001), *Co-5* on B7 (Fouilloux 1976; Campa et al. 2005), *Co-6* on B7 (Young and Kelly 1996a; Kelly et al. 2003), *Co-9* on B4 (Geffroy et al. 1999), and *Co-10* on B4 (Alzate-Marin et al. 2003). All these resistance loci segregate independently from one another except *Co-3* and *Co-9*, which are probably allelic (Méndez-Vigo et al. 2005).

Clusters of resistance genes to different pathogens or to different races of the same pathogen have been widely described in many plant species (Michelmore and Meyers 1998). In *P. vulgaris*, it has been reported the existence of genomic regions that concentrate resistance to several pathogens (Kelly et al. 2003). Recent studies have provided information on the genomic organization of regions harbouring resistance to anthracnose in common bean. These reports revealed the existence of families of resistance gene analogue sequences (RGAs) and/or resistance gene candidates arranged in clusters in the vicinity of *Co-2* (Geffroy et al. 1998; Creusot et al. 1999), *Co-4* (Melotto and Kelly 2001; Melotto et al. 2004), and *Co-3/Co-9* (Geffroy et al. 1999; Ferrier-Cana et al. 2003, 2005).

The main objectives of this work were the construction of a linkage map using the F₂ population derived from the cross between the two bean lines, Andecha and A252, and the characterization and mapping of the genes for specific resistance against six races of anthracnose present in these two lines. We discuss the possibility that clusters of different race-specific genes form most of the previously described anthracnose resistance loci.

Materials and methods

Plant material

Andecha is a very valuable large white seeded cultivar originating from a selection of landraces of Asturias

(Spain). A252 is a breeding line derived from the cross Carioca × Guanajuato 31, developed in CIAT (Cali, Colombia). To construct the linkage map, molecular marker analyses were carried out using DNA extracted from 85 F₂ plants from the cross Andecha × A252. Seventy-six F₃ families were used to characterize the corresponding F₂ plants for resistance to six races of anthracnose (races 6, 31, 38, 39, 65, and 357). The resistance to each race was independently evaluated in 10–40 plants per F₃ family. The 12 common bean differential cultivars (Pastor-Corrales 1991) were used to confirm the identity of the *C. lindemuthianum* isolates.

DNA extraction

Genomic DNA from the parental lines Andecha and A252 and from the F₂ progeny derived from the cross Andecha × A252, was isolated from lyophilized young leaves using the Nucleon™ PhytoPure™ Genomic DNA Extraction Kit (Amersham Biosciences) following the supplier's instructions.

PCR analysis

A total of 240 decamer primers obtained from Operon Technologies (Alameda, CA, USA) were used to screen the parents for polymorphism, and those generating repeatable and distinct fragments were selected. Polymerase chain reactions (PCRs) were performed in a 25- μ l total reaction mixture containing 30 ng of total DNA, 100 mM Tris-HCl, 100 mM KCl pH 8.3, 4 mM MgCl₂, 0.2 mM of each dNTP (Roche), 0.2 μ M of primer and 1.25 U Stoffel DNA polymerase (Applied Biosystems). Amplification reactions were performed in a PCR System 9600 (Applied Biosystems) programmed as follows: 94°C for 5 min, 35 cycles of 94°C for 35 s, 40°C for 2 min, and 72°C for 2 min with a ramp of 1 s per cycle, followed by a final step of 72°C for 7 min. The name of each RAPD marker is derived from an 'O' prefix for Operon primers, the letters identifying the Operon kit, Operon primer number, and the approximate size (bp) of the marker (subscript).

Polymerase chain reactions of the SCAR markers, BAC6 (Ariyaranthe et al. 1999), Phs (Kami et al. 1995), SAP6 (Miklas et al. 2000a), SAS13 (Young et al. 1998), SB8 (Rivkin et al. 1999), SB12 (Méndez-Vigo et al. 2005), SBD5 (Miklas et al. 2000b), ROC11 (Johnson et al. 1997), SI19 (Melotto and Kelly 1998), and SW13 (Melotto et al. 1996), were carried out as described by their corresponding authors. The amplification of SCAR-oli₅₀₀ was performed as described by Geffroy et al. (1998) modifying the annealing temperature (50°C instead of 58°C). The amplification of the SCAR SW12

(Miklas et al. 2000c) was carried out in a 25- μ l solution containing 25 ng of DNA, 100 mM Tris-HCl, 100 mM KCl pH 8.3, 5 mM MgCl₂, 0.2 mM of each dNTP (Roche), 0.2 μ M of each primer, and 1.25 U of Stoffel DNA polymerase (Applied Biosystems).

Polymerase chain reaction products amplified with RAPD, and SCAR primers were resolved on 2% agarose gels (except the SCAR Phs which was resolved on 8% polyacrylamide gels), stained with ethidium bromide and visualized under UV light. A 100-bp ladder (Amersham Biosciences) was used to measure the sizes of the amplification products.

RFLP analysis

RFLP analyses were performed with 33 probes designated as Bng, which are derived from a *Pst*I genomic library (Chase et al. 1991). These probes correspond to RFLP markers that were previously included in the University of Florida genetic map (Vallejos et al. 1992) and in the bean integrated linkage map (Freyre et al. 1998). The amplification products of the SCARs, SAS13 and SB8, were also used as probes for RFLP analysis after purification with GFX PCR DNA and Gel Band Purification Kit (Amersham Biosciences). Genomic DNA samples were digested with restriction enzymes *Dra*I, *Eco*RI, *Eco*RV, *Hind*III, *Taq*I, and *Hae*III (Amersham Biosciences). Restriction fragments were separated on 0.8% agarose gels and transferred to nylon membranes (Hybond-N; Amersham Biosciences) by vacuum blotting (VacuGene XL; Amersham Biosciences). Hybridizations of Southern blots with ³²P-labelled probes were carried out as described by Vallejos et al. (1992).

Seed colour (P locus)

The *P* locus controls the pigmentation on the seed coat; the dominant allele is present in A252 (coloured), and absent in Andecha (white). The segregation at this locus was followed by recording the presence or absence of pigmentation on the seeds formed by the F₂ plants.

Inoculation procedure and disease scoring

Six races of *C. lindemuthianum* were used: races 31, 39, 65, and 357, from the collection of the Crop and Soil Sciences Department (Michigan State University, US), and races 6 and 38 from the collection of SERIDA (Villaviciosa, Asturias, Spain). Isolates of each race were obtained from monosporic cultures maintained in fungus-colonized filter paper at -20°C for long-term

storage. The identity of each isolate was confirmed with the anthracnose differential set (Pastor-Corrales 1991). To obtain abundant sporulation, all races were grown at 19–21°C in darkness for about 10 days in Potato Dextrose Agar (Difco) (races 6, 31, 38, and 357) or in Marthur's agar (Marthur et al. 1950) (races 39 and 65). Spore suspensions were prepared by flooding the plates with 5 ml of 0.01% Tween 20 (Sigma) in sterile distilled water and scraping the surface of the culture with a spatula. Differential cultivars, parental lines, and F₃ families, were inoculated with a spore suspension of 1.2×10^6 spores/ml of the pathogen. Inoculations were carried out on 8 to 10-day-old seedlings in a climate chamber. The seedlings were sprayed with the aqueous conidial suspension and maintained at 20–22°C, 95–100% humidity, and 12 h photoperiod. The responses of the plants were evaluated after 7–9 days using a 1–9 scale where 1 is no visible symptoms and 9 very severely diseased or dead (Van Schoonhoven and Pastor-Corrales 1987).

Statistical analysis

Chi-square was used to test goodness-of-fit of observed to expected ratios in the F₂. The segregation analysis of the markers and the resistance genes was performed using MAPMAKER Macintosh version 2.0 (Lander et al. 1987). A LOD score of 3.0 was used as a linkage threshold with 0.25 as the maximum recombination fraction for linkage groups. Distances between ordered loci (cM) were calculated using the Kosambi mapping function.

Results

Construction of the genetic map

After screening the parental lines, Andecha and A252, 97 RAPD decamer primers were selected for their analysis in the F₂ population. These primers generated 164 dominant, and 8 codominant RAPD loci. Chi-square analyses indicated that the segregation of 11 of these RAPD markers (6.4%) deviated from the expected 3:1 ratio and were not included in the linkage analysis. All codominant RAPDs fitted the 1:2:1 expected segregation.

The Bng probes used constituted 28 codominant and 5 dominant RFLP markers that, in all cases, fitted the expected 1:2:1 and 3:1 segregations, respectively.

Ten out of the 12 SCARs analysed in the F₂ population segregated as dominant loci and fitted the 3:1 expectation. The remaining two SCARs, SAS13 (Young et al. 1998) and SB8 (Rivkin et al. 1999), produced no

differences in amplification pattern between the parental lines and did not segregate in the F_2 . The amplification products of these two SCARs were used as probes in RFLP analyses and segregations corresponding to codominant loci (1:2:1) were observed in both cases.

The seed colour trait (gene P,p) fitted a 3:1 monogenic dominant mode of inheritance.

The linkage analysis was performed on 85 F_2 individuals and 207 segregating markers (169 dominant and 38 codominant), using the multipoint linkage analysis software Mapmaker version 2.0 (Lander et al. 1987). The markers were sorted into distinct linkage groups using a LOD score of 3.0 as a linkage threshold with 0.25 as the maximum recombination fraction. This procedure resulted in the formation of 16 linkage groups with a number of markers ranging from 2 to 26, and 10 isolated markers (9 RAPDs and 1 Bng). The loci order of each linkage group was established following the full multipoint analysis mapping method. Later, correspondences of 16 of the linkage groups obtained with the 11 linkage groups of the bean integrated linkage map (Freyre et al. 1998) were established using Bng markers as anchor points. The resulting map (Fig. 1) includes 197 markers and covers 1,401.9 cM with an average marker distance of 7.1 cM.

Characterization of race-specific anthracnose resistance genes

Parental line Andecha was susceptible to races 6, 31, 38, 39, and 357 of *C. lindemuthianum* and showed an intermediate resistance phenotype to race 65, whereas parental line A252 was resistant to the six races.

For each of the races 6, 31, 38, 39, and 357, F_3 families were classified as having all individuals resistant (R), individuals resistant and susceptible (R/S), and all individuals susceptible (S). Table 1 shows the segregations of F_3 families for resistance to races 6, 31, 38, 39, and 357. Segregations for resistance to races 31 and 357 showed a good fit to the expected ratio for a single dominant gene (1 R:2 R/S:1 S), and were mapped on linkage groups B11 and B4, respectively (see Fig. 1). Segregations of resistance to races 6, 38, and 39 showed a good fit to the expected ratio for two independent dominant genes (7 R:8 R/S:1 S).

Table 2 shows the joint segregation of F_3 families for resistance to races 6, 31, 38, 39, and 357. Only 12 types of F_3 families (I–XII), considering their resistance spectra to these five races, were found. Types I–IX can be explained under the assumption of only two dominant genes being responsible for the different resistances, one of them conferring resistance to races 6, 38, 39, and 31 (located on B11) and the other conferring

resistance to races 6, 38, 39, and 357 (located on B4). However, types X, XI, and XII, indicate the occurrence of recombination between resistance specificities not supported by the two-single-genes hypothesis.

Table 3 shows the segregation of resistance to races 6, 38, 39, 31, and 357 within the six F_3 families constituting types X, XI, and XII. In these six families, the number of F_3 individuals scored for resistance to some races was increased in order to reduce the possibility of error.

The results indicated in Table 2 can be explained if different specific genes for resistance to races 6, 38, 39, 31, and 357, were arranged in two independent clusters on linkage groups B4 and B11. The B4 cluster would include the specific resistance genes R_1^6 , R_1^{38} , R_1^{39} , and R^{357} , and the B11 cluster would include the specific resistance genes R_2^6 , R_2^{38} , R_2^{39} , and R^{31} . Figure 2 illustrates this possibility.

Concerning resistance to race 65, all possible combinations of plants with total resistance (R_T), plants with intermediate resistance (R_I), and/or susceptible plants (S), were observed in the F_3 families. Table 4 shows the segregation of F_3 families for resistance to race 65, compared to the expected frequencies under the assumption of two independent dominant genes being responsible for the resistance to this race; one of them conferring total resistance in A252 (R_T^{65} , r_T^{65}), and the other conferring intermediate resistance in Andecha (R_I^{65} , r_I^{65}). Differences between the observed and expected frequencies are not significant.

The genotypes for gene R_T^{65} , r_T^{65} can be deduced from Table 4: F_3 families type I are homozygous resistant, $R_T^{65}R_T^{65}$, F_3 families types IV, V, and VII are heterozygous $R_T^{65}r_T^{65}$, and F_3 families types II, III, and VI are homozygous susceptible, $r_T^{65}r_T^{65}$. The segregation of this gene fits a 1:2:1 ratio ($\chi^2 = 3.98$; $P = 0.14$) and it was mapped at 10.2 cM (LOD = 13.99) of the R^{31} gene, in linkage group B11.

The genotypes for gene R_I^{65} , r_I^{65} can also be deduced in most F_3 families: types II and IV are homozygous for intermediate resistance, $R_I^{65}R_I^{65}$, types VI and VII are heterozygous $R_I^{65}r_I^{65}$, and types III and V are homozygous susceptible, $r_I^{65}r_I^{65}$. The segregation of this gene fitted the 1:2:1 ratio ($\chi^2 = 3.73$; $P = 0.15$) and it was mapped in linkage group B1.

Discussion

Localization of molecular markers linked to traits of interest

In regard to anthracnose resistance, the present map includes RAPD markers OF10₅₀₀ on LG B1 (probably




Fig. 1 Genetic linkage map of common bean in the F₂ population Andecha × A252. RAPD and SCAR markers, and race-specific resistance genes are indicated with designations described in the

text. *P* is a seed color gene. *Arrows* indicate common RFLP markers to the reference map published by Vallejos et al. (1992)

corresponding to OF10₅₃₀ linked to *Co-1*; Young and Kelly 1997), OQ04₁₄₀₀ on B11 (probably equivalent to OQ04₁₄₄₀ linked to *Co-2*; Young and Kelly 1996b), and OZ04₅₅₀ on B7, previously described as linked to *Co-6*

(Alzate-Marin et al. 1999). The amplification product of SCAreoli obtained in the present work (500 bp), was different from that described by Geffroy et al. (1998), but might identify close regions as both are

Table 1 Segregation for resistance to races 6, 31, 38, 39, and 357 of *Colletotrichum lindemuthianum* in F₃ Andecha × A252 families

Race	Observed frequency ^a			Expected frequency ^b			χ^2	Probability
	R	R/S	S	R	R/S	S		
6	28	39	2	30.2	34.5	4.3	0.745	0.39
31	20	32	15	16.8	33.5	16.8	0.881	0.64
38	32	42	2	33.3	38.0	4.8	0.468	0.49
39	29	38	2	30.2	34.5	4.3	0.402	0.53
357	16	35	17	17.0	34.0	17.0	0.088	0.96

^a F₃ families were classified as having all individuals resistant (R), individuals resistant and susceptible (R/S), and all individuals susceptible (S)

^b For races 6, 38, and 39 the expected ratio was 7:8:1 (two independent dominant genes) and for races 31 and 357 the expected ratio was 1:2:1 (one dominant gene)

Table 2 Joint segregation for resistance to races 6, 31, 38, 39, and 357 of *Colletotrichum lindemuthianum* in F₃ Andecha × A252 families

Type	Resistance spectrum of F ₃ families ^a						Frequency
	Race 6	Race 38	Race 39	Race 31	Race 357		
I	R/S	R/S	R/S	R/S	R/S		11
II	R/S	R/S	R/S	R/S	S		13
III	R/S	R/S	R/S	S	R/S		10
IV	R	R	R	R	R		10
V	R	R	R	R/S	R		2
VI	R	R	R	R	R/S		8
VII	R	R	R	R	S		2
VIII	R	R	R	S	R		3
IX	S	S	S	S	S		2
X	R/S	R/S	R	R/S	R/S		1
XI	R/S	R	R/S	R/S	R/S		3
XII	R	R	R	R/S	R/S		2

^a F₃ families were classified for each race as having all individuals resistant (R), individuals resistant and susceptible (R/S), and all individuals susceptible (S)

Table 3 Segregation of resistance to races 6, 38, 39, 31, and 357 within the six F₃ families proceeding from the cross Andecha × A252, corresponding to types X, XI, and XII of Table 2


Race	F ₃ families											
	F ₃ -75 (type X)		F ₃ -15 (type XI)		F ₃ -52 (type XI)		F ₃ -53 (type XI)		F ₃ -56 (type XII)		F ₃ -68 (type XII)	
	R	S	R	S	R	S	R	S	R	S	R	S
6	13	1	12	1	13	1	21	1	14	0	14	0
38	10	1	25	0	30	0	27	0	16	0	16	0
39	27	0	9	3	12	1	10	2	13	0	14	0
31	33	5	4	2	12	2	8	2	12	1	11	2
357	10	2	9	4	11	3	7	4	12	1	13	1

R resistant individuals, S susceptible individuals

located at the end of B11. SCAR markers SB12, SI19 and SW12, and RAPDs OAH18₁₁₀₀ and OY17₁₁₀₀ described as located in the vicinity of *Co-3/Co-9* resistance locus (Méndez-Vigo et al. 2005), were also mapped to B4 linkage group. Markers SI19 and OY17₁₁₀₀ had been previously described to be linked to the rust resistance gene *Ur-5* and SW12 linked to a QTL of resistance to bean golden yellow mosaic virus

(Miklas et al. 2002). The use of the amplification product of SAS13 (Young et al. 1998) as a probe in RFLP analysis (as polymorphism was not found between parents by direct amplification), allowed the indirect localization of *Co-4* locus on linkage group B8. This result agrees with the recent physical mapping of *Co-4* to chromosome 3 (which corresponds to linkage group B8; Melotto et al. 2004).

Fig. 2 An explanation for the results shown in Table 2, concerning the different F₃ family types observed (I–XII) for their combined resistance to races 6, 31, 38, 39, and 357. The genes proceeding from A252 providing resistance to these races are arranged in two clusters: the *B4* cluster includes the race-specific resistance genes R_1^6 , R_1^{38} , R_1^{39} , and R_1^{357} , and the *B11* cluster includes the race-specific resistance genes R_2^6 , R_2^{38} , R_2^{39} , and R_2^{31} . The corresponding *Andecha* haplotypes do not confer resistance to these races. *Types I–IX* are the different possible genotypes constituted by parental haplotypes. *Types X, XI, and XII* can only be explained with intra-cluster recombination. The relative positions of the different race-specific genes within each cluster is only tentative. The indicated positions explain types X–XII with a single recombination event



The present map also includes molecular markers linked to bean common mosaic virus resistance loci: SW13 (Melotto et al. 1996) linked to *I* gene on B2 linkage group, ROC11 (Johnson et al. 1997) linked to *bc-3* gene on B6 LG, and SBD5 linked to *bc-1³* (Miklas

et al. 2000b) on B3 LG. Two SCARs linked to common bacterial blight resistance were also mapped: SAP6 (Miklas et al. 2000a) on B10 and BAC6 on B2, although this location is not coincident with that given by Ariyaranthe et al. (1999) on B10.

Table 4 Segregation of F₃ Andecha × A252 families for resistance to race 65, compared to the expected frequencies under the assumption of two independent dominant genes being responsi-ble for the resistance to this race; one of them conferring total resistance (R_T⁶⁵, r_T⁶⁵), and the other conferring intermediate resistance (R_I⁶⁵, r_I⁶⁵)

F ₃ families			Genotypes of the corresponding F ₂ plants	Expected frequency
Type	Phenotypes ^a for resistance	Frequency		
I	R _T	22	R _T ⁶⁵ R _T ⁶⁵ –	15.8 (1/4)
II	R _I	5	r _T ⁶⁵ r _T ⁶⁵ R _I ⁶⁵ R _I ⁶⁵	3.9 (1/16)
III	S	6	r _T ⁶⁵ r _T ⁶⁵ r _I ⁶⁵ r _I ⁶⁵	3.9 (1/16)
IV	R _T /R _I	6	R _T ⁶⁵ r _T ⁶⁵ R _I ⁶⁵ r _I ⁶⁵	7.9 (1/8)
V	R _T /S	9	R _T ⁶⁵ r _T ⁶⁵ r _I ⁶⁵ r _I ⁶⁵	7.9 (1/8)
VI	R _I /S	6	r _T ⁶⁵ r _T ⁶⁵ R _I ⁶⁵ r _I ⁶⁵	7.9 (1/8)
VII	R _T /R _I /S	9	R _T ⁶⁵ r _T ⁶⁵ R _I ⁶⁵ r _I ⁶⁵	15.8 (1/4)

$$\chi^2 = 7.79 (P = 0.25)$$

^a F₃ families were classified as having all individuals with total resistance (R_T); all individuals with intermediate resistance (R_I); all individuals susceptible (S); individuals with total resistance and intermediate resistance (R_T/R_I); individuals with total resistance and susceptible (R_T/S); individuals with intermediate resistance and susceptible (R_I/S); individuals with total resistance, intermediate resistance and susceptible (R_T/R_I/S)

SB8 marker, a resistance analogue sequence, was located on linkage group B4 as described by Rivkin et al. (1999). The positions of genes *P* (seed colour) and *Phs* (Phaseolin: seed storage protein) on linkage group B7 agree with those of previous maps (Nodari et al. 1993; Freyre et al. 1998).

Characterization of race-specific anthracnose resistance genes

The intermediate resistance to race 65 present in Andecha can be explained by a single dominant gene (R_I⁶⁵) located on linkage group B1, linked to molecular markers Bng122 (17.6 cM) and OF10₅₀₀ (27.2 cM) previously described as linked to the *Co-1* gene (Young and Kelly 1997; Kelly et al. 2003). Among all characterized anthracnose resistance loci, *Co-1* is the only one originating in the Andean gene pool of common bean (Kelly and Vallejo 2004). As described before, land-race Andecha is considered to be Andean origin due to its morphological and biochemical characteristics.

The recombination between resistance specificities found in this work agrees with the assumption that total resistance to races 6, 31, 38, 39, 65, and 357, present in A252, is organized in two clusters. One of them, located on B4 linkage group, includes the race-specific resistance genes R_I⁶, R_I³⁸, R_I³⁹, and R³⁵⁷. The linkage between the R³⁵⁷ gene and the molecular markers OY17₁₁₀₀ (3.9 cM), SB12 (7.8 cM), OAH18₁₁₀₀ (19.2 cM), and SW12 (25 cM), indicates that this cluster corresponds to the *Co-3/Co-9* gene to which these markers were previously linked (Méndez-Vigo et al. 2005). The second cluster is located on linkage group B11 and includes the race-specific resistance genes R₂⁶, R₂³¹, R₂³⁸, R₂³⁹, and R_T⁶⁵. In this case,

the genetic distance between genes R³¹ and R_T⁶⁵ could be estimated as 10.2 cM. The linkage between the R³¹ gene and the molecular markers OQ04₁₄₀₀ (9.7 cM) and SCAreoli (14 cM), indicates that this cluster corresponds to the *Co-2* gene to which these markers were previously linked (Young and Kelly 1996b; Geffroy et al. 1998).

Genes providing resistance to different pathogens as well as to different races of the same pathogen have been reported to be clustered in chromosomal regions in many plant species (Crute and Pink 1996; Vear et al. 1997; Sharma et al. 2004). The results obtained in this work reveal that resistance genes *Co-2* and *Co-3/Co-9*, previously described as single major genes conferring resistance to several anthracnose races (Kelly and Vallejo 2004), should be considered as clusters of different genes conferring anthracnose race-specific resistance. The alleles of the *Co-3/Co-9* gene identified to date (Fouilloux 1976; Geffroy et al. 1999; Méndez-Vigo et al. 2005) could actually be haplotypes carrying different combinations of race-specific resistance genes. The tight linkage between the anthracnose resistance specificities *Co-y*, *Co-z*, and *Co-9* found by Geffroy et al. (1999) in the BAT93/JaloEEP558 RIL population, could also be interpreted as intra-cluster recombination of race-specific resistance genes. Also in the B4 cluster, recombination between race-specific genes for resistance to rust (Stavely 1984) and halo blight (Fourie et al. 2004) has been reported.

The *Co-2* locus had been only identified in the differential cultivar Cornell 49242 and to date no alleles had been reported (Kelly and Vallejo 2004). The *Co-2* haplotype present in A252 includes a specific gene for resistance to race 31 that would be absent at the *Co-2* cluster of Cornell 49242 (susceptible to race 31).

As found in other resistance loci in plants (Hulbert et al. 2001), the analysis of molecular organization of genomic DNA fragments corresponding to loci *Co-2* (Geffroy et al. 1998; Creusot et al. 1999), *Co-3/Co-9* (Geffroy et al. 1999; Ferrier-Cana et al. 2003, 2005), and *Co-4* (Melotto and Kelly 2001; Melotto et al. 2004) revealed the presence of clustered RGA sequences, suggesting a functional correspondence with the race-specific resistance genes described here.

Clusters of race-specific resistance genes could also constitute each of the remaining anthracnose resistance *Co-* loci of common bean. Segregation studies, using different anthracnose races could be carried out to determine whether it is possible to separate closely linked resistance specificities through recombination.

Acknowledgments This work was supported by grants AGL2001-2676-C02-01, RTA02-052-C2-2, and AGL2004-08145-C02-01 from the Ministerio de Ciencia y Tecnología, Spain. Astrid Pañeda and Belén Méndez-Vigo were recipients of salary fellowships from the Fundación para el Fomento de la Investigación Científica y Tecnológica, Principado de Asturias, Spain. Cristina Rodríguez-Suárez was recipient of a salary fellowship from the Ministerio de Ciencia y Tecnología, Spain. Races 31, 39, 65, and 357 were kindly provided by James D. Kelly (Michigan State University).

References

- Adam-Blondon A-F, Sévignac M, Bannerot H, Dron M (1994) SCAR, RAPD and RFLP markers linked to a dominant gene (*Are*) conferring resistance to anthracnose. *Theor Appl Genet* 88:865–870
- Alzate-Marin AL, Menarim H, de Carballo GA, de Paula Junior TJ, de Barros EG, Moreira MA (1999) Improved selection with newly RAPD markers linked to resistance gene to four pathotypes of *Colletotrichum lindemuthianum* in common bean. *Phytopathology* 89:281–285
- Alzate-Marin AL, Costa MR, Arruda KM, Gonçalves de Barros E, Moreira MA (2003) Characterization of the anthracnose resistance gene present in Ouro Negro (Honduras 35) common bean cultivar. *Euphytica* 133:165–169
- Aramuganathan K, Earle DE (1991) Nuclear DNA content of some important plant species. *Plant Mol Biol Rep* 9:208–218
- Ariyaranthe HM, Coyne DP, Jung G, Skroch PW, Vidaver AK, Steadman JR, Miklas PN, Basset MJ (1999) Molecular mapping of disease resistance genes for halo blight, common bacterial blight, and bean common mosaic virus in a segregating population of common bean. *J Am Soc Hortic Sci* 124:654–662
- Bannerot H (1965) Résultats de l'infection d'une collection de haricots par six races physiologiques d'anthracnose. *Ann Amélior Plant* 15:201–222
- Barrus MF (1915) An anthracnose-resistant red kidney bean. *Phytopathology* 5:303–311
- Blair MW, Pedraza F, Buendía HF, Gaitán-Solís E, Beebe SE, Gepts P, Tohme J (2003) Development of a genome-wide anchored microsatellite map for common bean (*Phaseolus vulgaris* L.). *Theor Appl Genet* 107:1362–1374
- Campa A, Rodríguez-Suárez C, Pañeda A, Giraldez R, Ferreira JJ (2005) The bean anthracnose resistance gene *Co-5*, is located in linkage group B7. *Annu Rep Bean Improv Coop* 48:68–69
- Chase CD, Ortega VM, Vallejos CE (1991) DNA restriction length fragments polymorphisms correlate with isozyme diversity in *Phaseolus vulgaris* L. *Theor Appl Genet* 81:806–811
- Creusot F, Macadré C, Ferrier Cana E, Riou C, Geffroy V, Sévignac M, Dron M, Langin T (1999) Cloning and molecular characterization of three members of the NBS-LRR subfamily located in the vicinity of the *Co-2* locus for anthracnose resistance in *Phaseolus vulgaris*. *Genome* 42:254–264
- Crute IR, Pink DAC (1996) Genetics and utilization of pathogen resistance in plants. *Plant Cell* 8:1747–1755
- Ferrier-Cana E, Geffroy V, Macadré C, Creusot F, Imbert-Bollere P, Sévignac M, Langin T (2003) Characterization of expressed NBS-LRR resistance gene candidates from common bean. *Theor Appl Genet* 106:251–261
- Ferrier-Cana E, Macadré C, Sévignac M, David P, Langin T, Geffroy V (2005) Distinct post-transcriptional modifications result into seven alternative transcripts of the CC-NBS-LRR gene *JAI1r* of *Phaseolus vulgaris*. *Theor Appl Genet* 110:895–905
- Fouilloux G (1976) L'anthracnose du haricot (*Colletotrichum lindemuthianum*, Sacc et Magn): nouvelles sources de résistance et nouvelles races physiologiques. *Ann Amélior Plant* 26(3):443–453
- Fourie D, Miklas P, Ariyaranthe H (2004) Genes conditioning halo blight resistance to races 1, 7, and 9 occur in a tight cluster. *Annu Rep Bean Improv Coop* 47:103–104
- Freyre R, Skroch PW, Geffroy V, Adam-Blondon AF, Shirmohamadali A, Johnson WC, Llaca V, Nodari RO, Pereira PA, Tsai SM, Tohme J, Dron M, Nienhuis J, Vallejos CE, Gepts P (1998) Towards an integrated linkage map of common bean. 4. Development of a core linkage map and alignment of RFLP maps. *Theor Appl Genet* 97:847–856
- Geffroy V, Creusot F, Falquet J, Sévignac M, Adam-Blondon A-F, Bannerot H, Gepts P, Dron M (1998) A family of LRR sequences in the vicinity of the *Co-2* locus for anthracnose resistance in *Phaseolus vulgaris* and its potential use in marker-assisted selection. *Theor Appl Genet* 96:494–502
- Geffroy V, Sicard D, de Oliveira JCF, Sévignac M, Cohen S, Gepts P, Neema C, Langin T, Dron M (1999) Identification of an ancestral resistance gene cluster involved in the coevolution process between *Phaseolus vulgaris* and its fungal pathogen *Colletotrichum lindemuthianum*. *Mol Plant Microbe Interact* 12(9):774–784
- Hulbert SH, Webb CA, Smith SM, Sun Q (2001) Resistance gene complexes: evolution and utilization. *Annu Rev Phytopathol* 39:285–312
- Johnson WC, Guzman P, Mandala D, Mkandawire ABC, Temple S, Gilbertson RL, Gepts P (1997) Molecular tagging of the *bc-3* for introgression into Andean common bean. *Crop Sci* 37:248–254
- Kami J, Velasquez VB, Debouck DG, Gepts P (1995) Identification of presumed ancestral DNA sequences of phaseolin in *Phaseolus vulgaris*. *Proc Natl Acad Sci USA* 92(4):1101–1104
- Kelly JD, Vallejo VA (2004) A comprehensive review of the major genes conditioning resistance to anthracnose in common bean. *HortScience* 39(6):1196–1207
- Kelly JD, Gepts P, Miklas PN, Coyne DP (2003) Tagging and mapping of genes and QTL and molecular marker-assisted selection for traits of economic importance in bean and cowpea. *Field Crops Res* 82:135–154

- Lander ES, Green P, Abrahamson J, Baarlow A, Daly MJ, Lincoln SE, Newburg L (1987) MAPMAKER: an interactive computing package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1:174–181
- Marthur RS, Barnett HL, Lilly VG (1950) Sporulation of *Colletotrichum lindemuthianum* in culture. *Phytopathology* 40:104–114
- Masterbroek C (1960) A breeding programme for resistance to anthracnose in dry shell haricot beans, based on a new gene. *Euphytica* 9:177–184
- Melotto M, Kelly JD (1998) SCAR markers linked to major disease resistance genes in common bean. *Annu Rep Bean Improv Coop* 41:64–65
- Melotto M, Kelly JD (2001) Fine mapping of the *Co-4* locus of common bean reveals a resistance gene candidate, *COK-4*, that encodes for a protein kinase. *Theor Appl Genet* 103:508–517
- Melotto M, Afanador L, Kelly JD (1996) Development of a SCAR marker linked to the *I* gene in common bean. *Genome* 39:1216–1219
- Melotto M, Coelho MF, Pedrosa-Harand A, Kelly JD, Camargo LEA (2004) The anthracnose resistance locus *Co-4* of common bean is located on chromosome 3 and contains putative disease resistance-related genes. *Theor Appl Genet* 109:690–699
- Méndez de Vigo B (2001) Mapa genético de *Phaseolus vulgaris* L. y resistencia a antracnosis en faba granja asturiana. Ph.D. thesis, University of Oviedo, Spain
- Méndez-Vigo B, Rodríguez-Suárez C, Pañeda A, Ferreira JJ, Giráldez R (2005) Molecular markers and allelic relationships of anthracnose resistance gene cluster B4 in common bean. *Euphytica* 141:237–245
- Michelmore RW, Meyers BC (1998) Clusters of resistance genes in plants evolve by divergent selection and birth and death process. *Genome Res* 8:1113–1130
- Miklas PN, Smith JR, Riley R, Grafton KF, Singh SP, Jung G, Coyne DP (2000a) Marker-assisted breeding for pyramided resistance to common bacterial blight in common bean. *Annu Rep Bean Improv Coop* 43:39–40
- Miklas PN, Larsen R, Victry K, Delorme R, Marma C, Riley RH, Kelly JD (2000b) Marker-assisted selection for the *bc-1²* gene for resistance to BCMV and BCMNV in common bean. *Euphytica* 116:211–219
- Miklas PN, Delorme R, Stone V, Daly MJ, Stavely JR, Steadman JR, Basset MJ, Beaver JS (2000c) Bacterial, fungal, virus disease loci mapped in a recombinant inbred common bean population (“Dorado/XAN 176”). *J Am Soc Hortic Sci* 125(2):476–481
- Miklas PN, Pastor-Corrales MA, Jung G, Coyne DP, Kelly JD, McClean PE, Gepts P (2002) Comprehensive linkage map of bean rust resistance genes. *Annu Rep Bean Improv Coop* 45:125–129
- Nodari RO, Tsai SM, Guzmán P, Gilbertson RL, Gepts P (1993) Towards an integrated linkage map of common bean III. Mapping genetic factors controlling host–bacteria interactions. *Genetics* 134:341–350
- Pastor-Corrales MA (1991) Estandarización de variedades diferenciales y designación de razas de *Colletotrichum lindemuthianum*. *Phytopathology* 81:694
- Pastor-Corrales MA, Tu JC (1994) Antracnosis. In: Pastor-Corrales MA, Schwartz HF (eds) Problemas de producción del frijol en los trópicos. Centro Internacional de Agricultura Tropical, Cali, Colombia, pp 87–120
- Pedrosa A, Vallejos CR, Bachmair A, Schweizer D (2003) Integration of common bean (*Phaseolus vulgaris* L.) linkage and chromosomal maps. *Theor Appl Genet* 106:205–212
- Rivkin MI, Vallejos CE, McClean PE (1999) Disease-resistance related sequences in common bean. *Genome* 42(1):41–47
- Schoonhoven A, Voysey O (1991) Common beans: research for crop improvement. Redwood, Melsham, Wiltshire, UK
- Sharma KD, Winter P, Kahl G, Muehlbauer J (2004) Molecular mapping of *Fusarium oxysporum* f. sp. *Ciceris* race 3 resistance gene in chickpea. *Theor Appl Genet* 108:1243–1248
- Stavely JR (1984) Genetics of resistance to *Uromyces phaseoli* in a *Phaseolus vulgaris* line resistant to most races of the pathogen. *Phytopathology* 74:339–344
- Vallejos EC, Sakiyama NS, Chase CD (1992) A molecular marker-based linkage map of *Phaseolus vulgaris* L. *Genetics* 131:733–740
- Van Schoonhoven A, Pastor-Corrales MA (1987) Standard system for the evaluation of bean germplasm. CIAT, Cali, Colombia
- Vear F, Gentzbittel L, Philippon J, Mouzeyar S, Mestries E, Roeckel-Drevet P, Tourvieille de Labrouhe D, Nicolas P (1997) The genetics of resistance to five races of downy mildew (*Plasmopara halstedii*) in sunflower (*Helianthus annuus* L.). *Theor Appl Genet* 95:584–589
- Young RA, Kelly JD (1996a) Characterization of the genetic resistance to *Colletotrichum lindemuthianum* in common bean differential cultivars. *Plant Dis* 80:650–654
- Young RA, Kelly JD (1996b) RAPD markers flanking the *Are* gene for anthracnose resistance in common bean. *J Am Soc Hortic Sci* 121:37–41
- Young RA, Kelly JD (1997) RAPD markers linked to three major anthracnose resistance genes in common bean. *Crop Sci* 37:940–946
- Young RA, Melotto M, Nodari RO, Kelly JD (1998) Marker assisted dissection of oligogenic anthracnose resistance in the common bean differential cultivar, G2333. *Theor Appl Genet* 96:87–94