

## Is the anthelmintic effect of heather supplementation to grazing goats always accompanied by anti-nutritional effects?

P. Frutos<sup>1†</sup>, J. Moreno-Gonzalo<sup>2</sup>, G. Hervás<sup>1</sup>, U. García<sup>3</sup>, L. M. M. Ferreira<sup>4</sup>, R. Celaya<sup>3</sup>, P. G. Toral<sup>1</sup>, L. M. Ortega-Mora<sup>2</sup>, I. Ferre<sup>2</sup> and K. Osoro<sup>3</sup>

<sup>1</sup>Estación Agrícola Experimental, Consejo Superior de Investigaciones Científicas (CSIC), Finca Marzanas, 24346 León, Spain; <sup>2</sup>SALUVET, Animal Health Department, Faculty of Veterinary Sciences, Complutense University of Madrid, Ciudad Universitaria s/n, 28040 Madrid, Spain; <sup>3</sup>Servicio Regional de Investigación y Desarrollo Agroalimentario (SERIDA), Área de Sistemas de Producción Animal, Consejería de Medio Ambiente y Desarrollo Rural, Principado de Asturias, 33300 Villaviciosa, Spain; <sup>4</sup>CECAV, Departamento de Zootecnia, Universidade de Trás-os-Montes e Alto Douro, PO Box 1013, 5000-911 Vila Real, Portugal

(Received 7 January 2008; Accepted 20 May 2008; First published online 22 July 2008)

To test the hypothesis that the beneficial anthelmintic effect of consuming moderate amounts of tannins may not always be accompanied by anti-nutritional effects in goats, two experiments were conducted. In the first, 48 Cashmere goats were randomly assigned to two treatments: supplementation with tannin-containing heather (6.4% total tannins) and non-supplementation. All goats grazed continuously from May to September under farm conditions in a mountainous area of northern Spain. The mean percentage of heather incorporated into the diet of the supplemented animals was 29.1%. Supplementation reduced the mean number of nematode eggs in faeces ( $P < 0.001$ ) and the goat mortality rate ( $P < 0.05$ ). The rumen ammonia concentration was markedly reduced in the goats receiving the heather supplement (160 v. 209 mg/l;  $P < 0.01$ ), while volatile fatty acid (VFA) concentrations were significantly greater (63.0 v. 53.6 mmol total VFAl;  $P < 0.05$ ). The heather-supplemented goats also showed a lower loss of live weight ( $P < 0.01$ ) and body condition score ( $P < 0.001$ ). In the second experiment, batch cultures of rumen microorganisms with rumen fluid from nine goats whose diet included 29% heather – or not, were used to incubate three substrates (pasture, pasture + heather and pasture + heather + polyethylene glycol) to investigate in vitro ruminal fermentation. Differences ( $P < 0.01$ ) among substrates were observed in terms of dry matter disappearance (DMD), in vitro true substrate digestibility (ivTSD), gas production and ammonia concentration, the greatest values always associated with the pasture substrate. Cultures involving rumen inoculum derived from goats receiving the heather-containing diet showed slightly lower DMD (46.9 v. 48.5 g/100 g;  $P < 0.05$ ), ivTSD (64.6 v. 65.9 g/100 g;  $P < 0.10$ ) and gas production (105 v. 118 ml/g;  $P < 0.001$ ) values, but much greater total VFA concentrations (48.5 v. 39.3 mmol/l;  $P < 0.05$ ), and suggest that the efficiency of ruminal fermentation in these animals was probably improved. Together, the results support the absence of a clear nutritional cost counteracting the beneficial anthelmintic effect of supplementing the diet of grazing goats with tannin-containing heather.

**Keywords:** body weight, gastrointestinal nematode, rumen fermentation, tannin

### Introduction

A growing number of studies relates the consumption of tannin-rich plants by small ruminants with the regulation of their gastrointestinal (GI) nematode populations. This could reduce dependence on conventional chemotherapy, to which many parasite species have developed resistance, and facilitate the sustainable control of GI nematode parasitism (Githiori *et al.*, 2006; Hoste *et al.*, 2006). Parasite-infected goats that consume moderate quantities of tannin-containing plants show improved resistance and resilience to parasites,

and significant decreases in faecal egg counts (FEC) and parasite burden (Min *et al.*, 2005; Paolini *et al.*, 2005; Osoro *et al.*, 2007a and 2007b). Goats can, however, also experience the anti-nutritional effects of tannins (e.g. Silanikove *et al.*, 1997; Ben Salem *et al.*, 2003).

It is widely accepted that the consumption of tannins by parasitized animals may result in either favourable or detrimental net effects depending on whether or not the positive anthelmintic action of these plant secondary metabolites outweighs their negative nutritional cost to the host (Houdijk and Athanasiadou, 2003). Nevertheless, some tannins have also been reported as capable of producing a positive nutritional effect when consumed in moderate

† E-mail: p.frutos@eae.csic.es

concentrations, which is associated with the reduction of dietary protein degradation in the rumen (Barry and McNabb, 1999; Mueller-Harvey, 2006).

Earlier studies (Osoro *et al.*, 2007a and 2007b) showed that grazing goats on a diet supplemented with tannin-containing heather had a lower FEC and a better performance, in comparison with animals on non-supplemented diets. On the basis of those studies, we hypothesized that the beneficial anthelmintic effect of the consumption of moderate amounts of tannins might not always be accompanied by marked anti-nutritional effects in goats. Two experiments were conducted to test this hypothesis. The first was performed under farm conditions in a less-favoured mountainous region of Spain where grazing animals are raised extensively and form part of the dynamics of the natural ecosystem. The second experiment involved an *in vitro* assay, with batch cultures of rumen microorganisms, to further investigate the nutritional effects of the inclusion of heather in the diet observed in the first.

## Material and methods

Experiments 1 and 2 were performed in accordance with Spanish Royal Decree 1201/2005 for the protection of animals used for experimental and other scientific purposes.

### Experiment 1 (*in vivo*)

#### Experimental site

This experiment was conducted in a mountainous area (altitude 1000 m) of northwestern Spain (6°53'W, 43°21'N; Sierra de San Isidro, Illano, Asturias) dominated by shrubby heather-gorse vegetation. Improved pastures were established in 2001 by removing the shrubs present and sowing perennial ryegrass (*Lolium perenne* L.) and white clover (*Trifolium repens* L.). A plot of approximately 20 000 m<sup>2</sup> previously managed homogeneously was divided into two identical plots of approximately 10 000 m<sup>2</sup> to accommodate all the animals used in this experiment.

During the experimental grazing season (May to September), the mean monthly rainfall ranged from 48 mm in August to 133 mm in September. The maximum and minimum temperatures were recorded in September (31°C) and May (2°C), respectively, the mean average monthly values ranging between 11.8°C in May and 17.8°C in July.

#### Animals and experimental design

Forty-eight lactating Cashmere goats (weighing 36.8 ± 1.13 kg at the beginning of the trial), balanced for live body weight (BW) and body condition score (BCS), were randomly assigned, together with their single kids, to one of two treatments: supplementation with heather (+H) and non-supplementation (-H). Each group was confined to one of the two plots of pasture (containing no heather, as explained above) of approximately 10 000 m<sup>2</sup> for the whole grazing season. The +H goats were offered, in the morning and once every 3 days, freshly cut heather *ad libitum*; the -H goats received no supplement. All animals were reared outdoors under farm conditions (i.e. not preventing nematode infection).

Two weeks before the experiment started, all animals were orally treated against GI nematodes with ivermectin (Oramec, Merial, Lyon, France) at about twice the recommended dose (2 mg/kg of BW). The animals were turned out on the pastures on 19 April and left grazing until 10 September.

#### Sampling procedures and analyses

**Pasture:** To monitor grass availability, the sward surface height was measured every 10 to 15 days using an HFRO swardstick (Barthram, 1986), taking 200 measurements at random in each plot. The botanical composition of the pasture was assessed in August using a point quadrat (Grant, 1981) and recording 200 vertical hits per plot. Pasture samples collected in August were analysed for dry matter (DM; ISO 6496:1999), organic matter (OM; ISO 5984:2002) and crude protein (CP; ISO 5983-2:2005). Neutral and acid detergent fibre (NDF and ADF) and acid detergent lignin (ADL) were determined by the method of Goering and Van Soest (1970) and Van Soest *et al.* (1991).

**Heather.** The species composition of the heather offered to the goats was assessed in August by recording 100 random contacts with the HFRO swardstick over the area where the heather was cut. Green shoots of this heather (less than 3 mm in diameter) were collected for chemical analysis. The nutritive quality (DM, OM, CP, NDF, ADF and ADL) was analysed following the same procedures as used to analyse the pasture. The total phenol (TP) and total tannin (TT) contents of the heather offered to the goats were determined using the Folin-Ciocalteu assay in combination with polyvinyl-polypyrrolidone, using tannic acid (Merck, Darmstadt, Germany) as the reference standard (Makkar, 2003).

**Diet selection.** In August, half of the +H animals were randomly chosen and subjected to estimations of the percentage of heather incorporated into the diet using *n*-alkane markers (Ferreira *et al.*, 2005). Samples of pasture, heather and faeces were collected and calculations performed using a least-squares optimization procedure, which minimizes the discrepancies between the actual concentration of alkanes in faeces (adjusted for incomplete faecal recoveries using the recovery values obtained by Ferreira *et al.* (2005)) and the estimated proportion in the diet.

**Parasitological procedures.** Approximately once per month between May and September, spot samples of goat faeces were individually collected by rectal grab to assess GI nematode egg excretion. The number of eggs per gram (epg) of faeces (FEC) was estimated using the modified McMaster technique (MAFF, 1978) with sodium chloride as the flotation medium, in which every egg is regarded as equivalent to 15 epg fresh faeces. Faecal cultures were performed for each group by pooling samples from the different individuals as described by MAFF (1978) and the genus of third-stage nematode larvae was identified according to Van Wyk *et al.* (2004).

**Body weight and body condition score.** All animals were weighed and their BCS assessed on a scale of 1 to 5 (1 = emaciated, 5 = fat; Russel, 1990), at the beginning and end of the experiment and at monthly intervals.

**Ruminal fermentation variables.** In August, after an overnight fast, a sample of ruminal fluid was individually collected from each goat via a stomach tube and visually checked to ensure that it did not contain saliva. Immediately afterwards, each fluid was strained through two layers of gauze and 4 ml was acidified with 4 ml 0.2 N HCl for ammonia determination. A further aliquot of 0.8 ml was added to 0.5 ml of a deproteinizing solution (2% metaphosphoric and 0.4% crotonic acids (wt/vol) in 0.5 N HCl) for determination of volatile fatty acid (VFA). All samples (two per animal and parameter to be analysed) were stored at  $-30^{\circ}\text{C}$  until analysis. The ammonia concentration was determined by colorimetry and VFA by gas chromatography, using crotonic acid as the internal standard, both in centrifuged samples (Frutos *et al.*, 2004).

**Mortality rate.** Goats were daily checked for clinical signs of illness. When an animal was considered terminally ill, it was humanely euthanized and complete *post mortem* examination was performed to discard mortality for other causes.

#### Statistical analysis

Sward height, FEC, BW and BCS data were analysed by repeated measures analysis using the MIXED procedure of the Statistical Analysis System program (SAS, 1999). The first measurement in May was used as the covariate. For FEC, BW and BCS, the animals were nested within the treatment to provide the error term to contrast the effect of heather supplementation. Data on FEC were log-transformed ( $\log_{10} x + 1$ ) to normalize their distribution.

The rumen variables were analysed by one-way analysis of variance (ANOVA), using the GLM procedure. The mortality rate was analysed using the  $\chi^2$  test (SAS, 1999).

Significant differences were declared at  $P < 0.05$  and tendencies at  $P < 0.10$ .

#### Experiment 2 (in vivo–in vitro)

##### Animals and diets

Nine non-lactating Cashmere goats ( $28.3 \pm 2.30$  kg BW) were used in this experiment, which was performed at an experimental research station. Animals were individually fed a mixture of alfalfa and grass hays (AGH; 18 g DM/kg of BW) once a day over a 14-day period. They were then switched to a diet containing 71% of the same mixture of hays and 29% heather (AGH + H; 18 g DM/kg of BW) for 12 more days. The heather was collected from the same area as in Experiment 1. These diets (treatments) were designed on the basis of the outcome of diet selection (see results from Experiment 1), to experimentally simulate the –H and +H treatments.

Table 1 shows the chemical composition of the mixture of grass and alfalfa hays and of the heather used in this experiment.

**Table 1** Chemical composition (g/kg of DM, except for DM itself (g/kg of fresh matter)) of feeds and diets used in Experiments 1 and 2

	DM	OM	CP	NDF	ADF	ADL	TT <sup>1</sup>	TP <sup>1</sup>
Experiment 1 <sup>2</sup>								
Pasture	328	949	143	590	269	25	nd	nd
Heather	423	979	61	557	436	224	64	94
Experiment 2 (diets) <sup>3</sup>								
AGH	865	899	154	591	347	61	nd	nd
AGH + H	776	921	127	581	364	101	23	33

DM = dry matter; OM = organic matter; CP = crude protein; NDF = neutral detergent fibre; ADF = acid detergent fibre; ADL = acid detergent lignin; TT = total tannins; TP = total phenols; nd = not determined.

<sup>1</sup>g tannic acid equivalents/kg of DM.

<sup>2</sup>Samples collected in August.

<sup>3</sup>AGH = alfalfa and grass hays; AGH + H = alfalfa and grass hays (71%) + heather (29%).

#### In vitro ruminal fermentation

The effect of the diet on *in vitro* ruminal fermentation was studied in four runs of incubation using a modification of the gas production technique described by Theodorou *et al.* (1994). On day 12 of the experiment, after an overnight fast, ruminal fluid was collected from each animal by stomach tube (as in Experiment 1). The collected samples were transferred to the laboratory in pre-warmed thermos flasks, strained through a double layer of muslin, and kept under  $\text{CO}_2$  flushing. The nine inocula were mixed to form a single inoculum (run 1). The whole process was repeated on day 14 to obtain the replicate (run 2). When the goats received the heather-containing diet, rumen fluids were again withdrawn on days 24 (run 3) and 26 (run 4).

Samples of the pasture and heather collected in August during Experiment 1, ground through a 1-mm screen, were used to provide two substrates: pasture (P; 100% pasture) and pasture + heather (P + H; 71% pasture and 29% heather). A third substrate (P + H + PEG) was prepared by adding 1 g per flask of a tannin-binding agent (polyethylene glycol, PEG, MW 6000; Fluka Chemie GmbH, Buchs, Switzerland) – which inhibits the effect of tannins (Makkar, 2003) – to the P+H substrate. The large quantity of PEG was chosen because the efficiency of this agent is believed to depend on the tannins/PEG ratio, and to avoid the possibility that too small an amount of this inert compound might be associated with a lack of effect.

Each incubation run included a total of twelve 125 ml sealed serum flasks. Three samples of each substrate ( $\approx 500$  mg) and three blanks (i.e. buffered rumen fluid without substrate) were incubated at  $39^{\circ}\text{C}$  with 10 ml rumen fluid and 40 ml phosphate–bicarbonate buffer. The buffer solution was prepared as described by Goering and Van Soest (1970), with the exception that no trypticase was added.

Gas production was determined by measuring headspace gas pressure at 24 h post inoculation. Pressure values, corrected for the quantity of substrate OM incubated and gas released from the blanks, were used to generate gas

volume estimates using a predictive equation derived from earlier simultaneous pressure and volume measurements (Hervás *et al.*, 2005). Immediately afterwards, the fermentations were ended by swirling the bottles on ice. The samples were then centrifuged at  $976 \times g$  for 10 min and aliquots of the supernatant collected for ammonia and VFA determinations as in Experiment 1.

*In vitro* DM disappearance (DMD; g/kg) and *in vitro* true substrate digestibility (ivTSD) after 24 h of incubation were estimated by filtering the residues using pre-weighed sintered glass crucibles (100 to 60  $\mu\text{m}$ ; Pyrex, Stone, UK) and determining the NDF content, as reported in Frutos *et al.* (2004).

**Chemical analyses**

All chemical analyses were performed as in Experiment 1.

**Statistical analysis**

Data for *in vitro* ruminal variables were analysed by one-way ANOVA, using the GLM procedure of the Statistical Analysis System program (SAS, 1999) to examine the effects of the two main sources of variation, goat diets and substrates, as well as their interaction. The mean value of each set of three flasks per substrate and run was considered as the experimental unit. Significance was set at  $P < 0.05$ ; trends were declared at  $P < 0.10$ .

**Results**

**Experiment 1**

*Sward height, and pasture and heather botanical and chemical composition*

As shown in Figure 1, the available sward height differed significantly between treatments, with greater values being recorded in the pasture used by the +H animals ( $P < 0.001$ ). Mean sward height gradually fell from 10.2 cm at the beginning of the experiment (May) to 3.7 cm at the end (September) ( $P < 0.001$ ).

The botanical composition of the available pasture in August was on average 6.8% white clover, 21.9% perennial ryegrass, 43.1% other grasses (mainly *Agrostis capillaris* L.) and 27.9% dead matter, with no differences between treatments. The chemical composition of the pasture samples taken in August showed a CP content of 143 g/kg of DM, and values for NDF, ADF and ADL of 590, 269 and 25 g/kg of DM, respectively (Table 1).

The heather offered was mostly composed of *Calluna vulgaris* (L.) Hull (61%), with 25% *Erica umbellata* L., 12% *E. cinerea* L., 1% *E. tetralix* L. and 1% *Daboecia cantabrica* (Hudson) C. Koch. Chemical composition analysis showed a lower CP content (61 g/kg of DM) and greater cell wall content (557, 436 and 224 g/kg of DM for NDF, ADF and ADL, respectively), as expected for shrub species. TP and TT contents were estimated to be 94 and 64 g tannic acid equivalents/kg of DM, respectively (Table 1).


**Diet composition**

The mean percentage of heather incorporated into the diet of the +H animals was 29.1% ( $\pm 2.06\%$ ) in August. The remaining 70.9% corresponded to herbaceous species.

**Faecal egg counts and goat mortality**

Supplementation with heather reduced the mean FEC ( $P < 0.001$ ; see Figure 1); values ranged from 15 epg at the beginning of the trial to a maximum of 2653 epg at the end of August. The -H goats with the greatest FEC in the two samplings performed in August died before the September sampling (seven in total), which explains the reduction seen in Figure 1. Supplementation with heather significantly reduced the mortality rate ( $P < 0.05$ ): at the end of the grazing period, two +H and eight -H does had to be euthanized.

The nematode genera identified were *Trichostrongylus*, *Teladorsagia* and *Oesophagostomum*. *Trichostrongylus* was the predominant genus (60% to 95%) over the entire experimental period; *Teladorsagia* and *Oesophagostomum* were detected in lower percentages (5% to 25%).



**Figure 1** Faecal gastrointestinal nematode egg counts (FEC; eggs/g; solid lines —) and sward height (cm; short-dash lines - -) over the experimental season in grazing goats whose diets were supplemented – or not – with heather. Mean log-transformed FEC values = 2.20 v. 2.66 eggs/g for supplemented v. non-supplemented animals (s.e.d. = 0.115;  $P < 0.001$ ), and mean sward heights = 6.8 v. 5.7 cm for supplemented v. non-supplemented animals (s.e.d. = 0.07;  $P < 0.001$ ) (s.e.d. = standard error of the differences of means).

**Table 2** Body weight (BW, kg) and body condition score (BCS) during the experimental grazing period (from May to September) in goats supplemented (+H) or not (-H) with heather

	BW			BCS		
	-H	+H	s.e.	-H	+H	s.e.
May	37.0	36.3	1.12	2.57	2.56	0.042
June	34.5	34.2	0.89	2.57	2.59	0.043
July	36.2 <sup>b</sup>	37.7 <sup>a</sup>	0.92	2.53 <sup>b</sup>	2.85 <sup>a</sup>	0.049
August	30.3 <sup>b</sup>	32.4 <sup>a</sup>	0.83	2.17 <sup>b</sup>	2.38 <sup>a</sup>	0.068
September	29.7 <sup>b</sup>	30.7 <sup>a</sup>	0.87	1.45 <sup>b</sup>	2.00 <sup>a</sup>	0.076
<i>P</i> value <sup>1</sup>						
Treatment	<0.01			<0.01		
Date of sampling	<0.01			<0.01		
Treatment × date	<0.01			<0.01		

s.e. = standard error of the mean.

<sup>1</sup>Probability value.

<sup>a,b</sup>For each variable, means in a row with a different superscript differ significantly ( $P < 0.05$ ).

### Live weight and body condition changes

Table 2 shows that heather supplementation was responsible for the smaller losses of live BW ( $P < 0.01$ ) and BCS ( $P < 0.01$ ) recorded over the experimental grazing season in the +H animals.

### Ruminal fermentation variables

Table 3 shows the ammonia and VFA concentrations in the ruminal fluid of the experimental animals. The rumen ammonia concentration was markedly reduced in the goats receiving tannin-containing heather ( $P < 0.01$ ). On the contrary, total VFA concentration was significantly greater in these animals ( $P < 0.05$ ), mainly due to an increase in acetic acid (approximately 26%) but also in propionic acid (approximately 16%). The concentrations of valerate and branched VFAs (referred to as 'others' in Table 3), however, decreased. The molar proportions of VFA (data not shown) were also modified by supplementation with heather ( $P < 0.001$ ), with increases in acetate and reductions in butyrate, valerate and branched VFAs.

### Experiment 2

As shown in Table 4, significant differences in gas production, DMD, ivTSD, and ammonia and VFA concentrations were observed due to both goat diet (i.e. including or not heather: AGH v. AGH + H) and substrates incubated. However, no significant interaction (diet × substrate) was found for any *in vitro* rumen fermentation variables ( $P > 0.10$ ).

Cultures with rumen inoculum derived from goats receiving the heather-containing diet (AGH + H) showed slightly lower values of DMD (-3%), ivTSD (-2%) and gas production (-11%) compared with cultures from AGH animals. On the contrary, those incubations (AGH + H) returned greater values of total VFA (23%), mainly accounted for by greater concentrations of acetate (34%) and butyrate (21%). The molar proportions of VFA (data not shown) tended to be modified by the incorporation of

**Table 3** Ammonia (mg/l) and volatile fatty acid (VFA, mmol/l) concentrations in the ruminal fluid of grazing goats whose diets were supplemented (+H) or not supplemented (-H) with heather

	-H	+H	s.e.	<i>P</i> value <sup>1</sup>
Ammonia	209.4 <sup>a</sup>	159.7 <sup>b</sup>	9.95	<0.01
Total VFA	53.61 <sup>b</sup>	62.98 <sup>a</sup>	2.340	0.02
Acetate	36.11 <sup>b</sup>	45.47 <sup>a</sup>	1.845	<0.01
Propionate	8.80 <sup>b</sup>	10.25 <sup>a</sup>	0.500	0.07
Butyrate	5.48	5.08	0.270	0.34
Others <sup>2</sup>	3.22 <sup>a</sup>	2.17 <sup>b</sup>	0.137	<0.01

s.e. = standard error of the mean.

<sup>1</sup>Probability value.

<sup>2</sup>Calculated as the sum of valerate, isovalerate, isobutyrate and caproate.

<sup>a,b</sup>Means in a row with a different superscript differ significantly ( $P < 0.05$ ).

heather into the diet ( $P < 0.10$ ), with increases in acetate and reductions in propionate, and valerate and branched VFAs (referred to as 'others' in Table 4).

Differences among substrates ( $P < 0.01$ ) were observed in terms of gas production, DMD, ivTSD and ammonia concentrations, the greatest values always being found for the P substrate and the lowest for the P + H substrate. When PEG was added to the mixture (P + H + PEG), gas production and ivTSD were significantly increased. However, the addition of PEG did not significantly inhibit the negative effect of heather on DMD and ammonia concentration.

No significant differences among substrates were observed for the concentration of any particular VFA.

### Discussion

The control of GI parasites with anthelmintics is becoming increasingly difficult worldwide due to drug resistance and concerns over pesticide residues in the food chain and the general environment (Van Houtert and Sykes, 1996). Thus, the greater variety of control methods used in combination, the longer we can expect to have effective worm control. The use of tannin-containing plants has proven effective in the control of GI parasite infections, as confirmed in this study where supplementing the diets of grazing goats with tannin-containing heather significantly reduced FEC (see Figure 1). This is in line with previous reports for the same area by Osoro *et al.* (2007a and 2007b). However, the mean FEC values (as high as 2652 epg) were notably different from those found in the above papers, which is almost certainly related to the effect of climatic conditions on parasite infections (Stromberg, 1997).

Several authors have reported that the intake of low-to-moderate quantities of condensed tannins (CTs) may have beneficial effects on parasitized ruminants. For example, we reported reductions in FEC of up to 75%, compared with control goats, in an experiment that included supplementation with heather containing between 7% and 8.6% tannin (expressed in tannic acid equivalents; Osoro *et al.*, 2007b). Butter *et al.* (2000) observed a reduction in FEC when 5% of a commercial extract of quebracho (*Schinopsis* spp.)

**Table 4** Gas production (ml/g OM), dry matter disappearance (DMD; g/100 g), in vitro true substrate digestibility (ivTSD; g/100 g), ammonia (mg/l) and VFA concentration (mmol/l) in batch cultures after 24 h incubation

	Goat diet <sup>1</sup>			Substrate <sup>2</sup>				P value <sup>3</sup>		
	AGH	AGH + H	s.e.	P	P + H	P + H + PEG	s.e.	Diet	Substrate	Diet × subst
Gas production	118.8 <sup>a</sup>	105.2 <sup>b</sup>	1.55	123.5 <sup>a</sup>	102.1 <sup>c</sup>	110.4 <sup>b</sup>	1.903	<0.01	<0.01	0.39
DMD	48.48 <sup>a</sup>	46.94 <sup>b</sup>	0.358	52.82 <sup>a</sup>	44.40 <sup>b</sup>	45.90 <sup>b</sup>	0.439	0.03	<0.01	0.55
ivTSD	65.95 <sup>a</sup>	64.61 <sup>b</sup>	0.431	67.23 <sup>a</sup>	63.08 <sup>b</sup>	65.54 <sup>a</sup>	0.775	0.07	<0.01	0.82
Ammonia	287.7	284.7	3.51	308.6 <sup>a</sup>	269.4 <sup>b</sup>	280.74 <sup>b</sup>	4.30	0.57	<0.01	0.31
Total VFA	39.27 <sup>b</sup>	48.45 <sup>a</sup>	1.830	45.44	41.79	44.34	2.241	0.02	0.54	0.97
Acetate	23.65 <sup>b</sup>	31.81 <sup>a</sup>	1.333	28.91	26.83	27.43	1.632	<0.01	0.67	0.82
Propionate	10.44	10.82	0.746	10.70	9.99	11.22	0.913	0.74	0.66	0.81
Butyrate	3.01 <sup>b</sup>	3.65 <sup>a</sup>	0.208	3.56	3.10	3.33	0.255	0.08	0.48	0.97
Others <sup>4</sup>	2.17	2.18	0.126	2.28	1.88	2.36	0.155	0.97	0.15	0.88

s.e. = standard error of the mean.

<sup>1</sup>AGH = alfalfa and grass hays; AGH + H = alfalfa and grass hays (71%) + heather (29%).

<sup>2</sup>P = pasture; P + H = pasture (71%) + heather (29%); PEG = polyethylene glycol.

<sup>3</sup>Probability value.

<sup>4</sup>Calculated as the sum of valerate, isovalerate, isobutyrate and caproate.

<sup>a,b,c</sup>For each source of variation (goat diet or substrate), means in a row with a different superscript differ significantly ( $P < 0.05$ ).

CTs was included in a low protein diet. Min *et al.* (2005) reported a great reduction in FEC when does were fed *Sericea lespedeza* (*Lespedeza cuneata*) containing 15.2% tannins (using an internal standard from the plant itself as the reference).

In the present study, FEC reduction was achieved with the dietary incorporation of an estimated 29% heather containing 6.4% tannin (which may be considered a low-to-moderate intake of tannins). The lowest dietary tannin threshold needed for anthelmintic effects remains unclear, however. Moreover, it should be remembered that different plants contain tannins with different structures and hence reactivities, so the notion of a 'low-to-moderate' content could be misleading. In addition, different research groups use different methods to analyse this large group of phenolic compounds, but the equivalents in which the tannin content are expressed (e.g. tannic acid, internal standards extracted from the plants themselves, etc.) are not always given, which means that comparisons cannot be made (Makkar, 2003; Álvarez del Pino *et al.*, 2005).

A number of investigations have focused on the potential use of CTs to control GI parasitism because these are relatively stable in the digestive tract of ruminants and rarely have toxic effects (Mueller-Harvey, 2006). CTs may affect protein nutrition by reducing dietary protein degradation in the rumen, thus enhancing the amount of protein available for digestion in the small intestine (Barry and McNabb, 1999; Frutos *et al.*, 2000; Mueller-Harvey, 2006). According to Coop and Kyriazakis (2001), protein nutrition, and therefore CTs, may influence the development and consequences of parasitism by increasing the resilience and resistance of the host or by directly affecting the parasite populations.

Given the low content of protein in the offered heather (61 g/kg of DM), it seems very unlikely that the anthelmintic effect of its supplementation would be due to a greater

availability of protein, although there might be a greater availability of non-degradable protein due to the effect of the tannin (Barry and McNabb, 1999). A common characteristic of GI parasitism is an increased loss of endogenous protein to the GI tract (Coop and Kyriazakis, 2001) and some authors have suggested that the inclusion of CTs might be an alternative to providing high protein rations as a means of reducing the consequences of parasitism (Van Houtert and Sykes, 1996; Butter *et al.*, 2000; Coop and Kyriazakis, 2001).

The supplementation with tannin-containing heather significantly reduced the goat mortality rate, which is consistent with previous studies reporting the positive effect of tannin consumption on the resilience of goats (Paolini *et al.*, 2005).

In addition to the effect of heather supplementation and due to the differences observed between treatments in the sward height, it cannot be completely ruled out that the -H goats consumed more contaminated pasture than +H goats. A shorter sward may have larger parasite burdens because it is easier for the parasites to migrate to the top (Niezen *et al.*, 1998), so the non-supplemented animals might be at a higher risk of ingestion of nematode infective larvae. Nevertheless, as both experimental plots had the same previous management, they presumably had a similar parasitic status at the start of the trial.

As observed in earlier work (Osoro *et al.*, 2007a and 2007b), the negative BW gains agree with the relationship established between green sward height and BW change (Merchant and Riach, 1994). Considerable BW losses were observed after July, when the sward height fell below 6 to 7 cm (see Figure 1), probably due to the limited availability of pasture and the increase in parasite burden during the grazing season (Coop and Kyriazakis, 2001). However, losses were mitigated by the effect of heather intake. Houdijk and Athanasiadou (2003) reported evidence suggesting

that the consumption of CT-rich plants can result in reduced parasitism and improved performance in parasitized small ruminants, which would probably be the outcome of indirect nutritional benefits (e.g. extra protein availability for the parasitized grazing hosts). On the other hand, those authors also suggest that the net benefit in the performance of parasitized hosts might be the result of the anti-parasitic effects of CTs outweighing their anti-nutritional effects.

The performance results found in this study were supported by the nutritional outcomes. Although the results on ruminal fermentation obtained *in vitro* and *in vivo* were not exactly the same, as expected, they followed a very similar pattern and support the general conclusion that the dose of tannins ingested with the heather diet was insufficient to adversely affect the majority of the rumen microbial populations in goats adapted to its consumption.

Several authors have indicated that dietary concentrations of <50 g CT/kg DM are nutritionally beneficial (these recommendations originated mainly from feeding trials with *Lotus* species and cannot be directly applied to other feeds; Mueller-Harvey, 2006). However, erroneous generalizations have persisted since the first reviews in the 1960s and 1970s stated that tannins are harmful or toxic to mammals (Mueller-Harvey, 2006). The main benefit of tannins in ruminant nutrition stems from their negative effect on proteolysis and the first consequence is a lower concentration of ammonia in the ruminal fluid, as observed in our *in vivo* and *in vitro* approaches. The differences between assays 1 and 2 may be due to the fact that, in *in vitro* studies, lower rumen fluid ammonia concentrations can only be attributed to lower concentrations of protein in the substrate, a reduced deamination of amino acids or a greater rate of uptake by bacteria. In addition, differences may also be explained because in the *in vitro* experiment 29% of the pasture was replaced by exactly the same amount of heather (while substitution in the *in vivo* grazing trial was probably incomplete), and because the response of the ruminal microorganisms to tannins may not be exactly the same *in vivo* and *in vitro*. The fact that no significant differences in ammonia concentration were observed between substrates P + H and P + H + PEG suggests that other factors, apart from tannins, may also be at work (e.g. the lower nutritive value of P + H compared with P). In the grazing experiment, a significant and negative correlation was found between the percentage of heather incorporated into the diet and the rumen ammonia concentration ( $r = -0.60$ ;  $P < 0.05$ ).

As is well known, VFAs are the most important by-products of fermentation and the major energy source of ruminants. The higher concentrations of VFAs for the supplemented goats in both trials (17% and 23%, respectively) are in line with the better performance (BW and BCS) of these animals. The fact that no significant differences were found in total VFA concentration between substrates, but were found when comparing goat diets, suggests an adaptation of rumen microorganisms to the consumption of these phenolic compounds (Smith *et al.*, 2005) and a subsequently better digestive utilization of the diet.

The direct effects reported for tannins on rumen VFAs are not consistent and different authors have indicated different results (e.g. Makkar *et al.*, 1995; Hervás *et al.*, 2003); these discrepancies probably depend on the amount and type of tannins and on the animal species that ingest them. Results from Experiments 1 and 2 revealed a significant increase in acetate production when heather was present, as well as differences in the fermentation pathways, reflected in the different molar proportions. The increase in acetate and the reduction in gas production may be accounted for by an effect of tannins on acetate-utilizing rumen bacteria, which is probably related to the beneficial and known inhibition of methane-producing microorganisms by these secondary compounds (Tavendale *et al.*, 2005). These results support the improved efficiency of ruminal fermentation in animals consuming tannin-containing heather and confirm those reported in a previous grazing experiment performed under similar conditions (Osoro *et al.*, 2007b).

Comparisons of the results for goats adapted *v.* non-adapted to the consumption of heather (AGH *v.* AGH + H; Table 4) show the latter to be associated with reductions in gas production, DMD and iVTD (in accordance with that widely reported for diets containing tannins; Frutos *et al.*, 2004; Mueller-Harvey, 2006) of a lower magnitude than the improvements in the mean indicators for the utilization of nitrogen or energy (ammonia and VFA concentrations).

In conclusion, the present results suggest there is no substantial nutritional cost counteracting the anthelmintic effect of supplementing the diets of grazing goats with heather, and they therefore support the hypothesis under test. Further research, including the estimation of total pasture and heather intake, would help explain the effects of heather supplementation on nutrition and performance in grazing goats. Eventually, it would also contribute to the development of a sustainable method for controlling GI nematode parasitism in extensive goat production systems.

### Acknowledgements

The authors would like to thank the Carbayal Research Station staff for their help in the field work. This work was supported by the Spanish National Institute for Agrarian and Food Research and Technology (INIA, Project RTA2007-00098).

### References

- Álvarez del Pino MC, Hervás G, Mantecón AR, Giráldez FJ and Frutos P 2005. Comparison of biological and chemical methods, and internal and external standards, for assaying tannins in Spanish shrub species. *Journal of the Science of Food and Agriculture* 85, 583–590.
- Barry TN and McNabb WC 1999. The implications of condensed tannins on the nutritive value of temperate forages fed to ruminants. *British Journal of Nutrition* 81, 263–272.
- Barthram GT 1986. Experimental techniques: the HFRO swardstick. The Hill Farming Research Organisation Biennial Report 1984–85, pp. 29–30. HFRO, Bush Estate, Penicuik, Midlothian, UK.
- Ben Salem H, Ben Salem I, Nefzaoui A and Ben Saïd MS 2003. Effect of PEG and olive cake feed blocks supply on feed intake, digestion, and health of goats given kermes oak (*Quercus coccifera* L.) foliage. *Animal Feed Science and Technology* 110, 45–59.

- Butter NL, Dawson JM, Wakelin D and Buttery PJ 2000. Effect of dietary tannin and protein concentration on nematode infection (*Trichostrongylus colubriformis*) in lambs. *Journal of Agricultural Science* 134, 89–99.
- Coop RL and Kyriazakis I 2001. Influence of host nutrition on the development and consequences of nematode parasitism in ruminants. *Trends in Parasitology* 17, 325–330.
- Ferreira LMM, Oliván M, García U, Rodrigues MAM and Osoro K 2005. Validation of the alkane technique to estimate diet selection of goats grazing heather-gorse vegetation communities. *Journal of the Science of Food and Agriculture* 85, 1636–1646.
- Frutos P, Hervás G, Giráldez FJ, Fernández M and Mantecón AR 2000. Digestive utilization of quebracho-treated soya bean meal in sheep. *Journal of Agricultural Science* 134, 101–108.
- Frutos P, Hervás G, Giráldez FJ and Mantecón AR 2004. An *in vitro* study on the ability of polyethylene glycol to inhibit the effect of quebracho tannins and tannic acid on rumen fermentation in sheep, goats, cows and deer. *Australian Journal of Agricultural Research* 55, 1125–1132.
- Githiori JB, Athanasiadou S and Thamsborg SM 2006. Use of plants in novel approaches for control of gastrointestinal helminths in livestock with emphasis on small ruminants. *Veterinary Parasitology* 139, 308–320.
- Goering MK and Van Soest PJ 1970. Forage fiber analysis (apparatus, reagents, procedures and some applications). *Agricultural handbook no. 379*. ARS-USDA, Washington, DC, USA.
- Grant SA 1981. Sward components. In *Sward measurement handbook* (ed. J Hodgson, RD Baker, A Davies, AS Laidlaw and JD Leaver), pp. 71–92. British Grassland Society, Hurley, Maidenhead, Berkshire, UK.
- Hervás G, Frutos P, Giráldez FJ, Mantecón AR and Álvarez del Pino MC 2003. Effect of different doses of quebracho tannins extract on rumen fermentation in ewes. *Animal Feed Science and Technology* 109, 65–78.
- Hervás G, Frutos P, Giráldez FJ, Mora MJ, Fernández B and Mantecón AR 2005. Effect of preservation on fermentative activity of rumen fluid inoculum for *in vitro* gas production techniques. *Animal Feed Science and Technology* 123–124, 107–118.
- Hoste H, Jackson F, Athanasiadou S, Thamsborg SM and Hoskin SO 2006. The effects of tannin-rich plants on parasitic nematodes in ruminants. *Trends in Parasitology* 22, 253–261.
- Houdijk JGM and Athanasiadou S 2003. Direct and indirect effects of host nutrition on ruminant gastrointestinal nematodes. In *Matching herbivore nutrition to ecosystems biodiversity*. VI International Symposium on Nutrition of Herbivores (ed. L 't Mannetje, L Ramírez, C Sandoval-Castro and JC Ku-Vera), pp. 213–236. Univ. Aut. Yucatán, Mérida, México.
- MAFF 1978. *Manual of veterinary parasitological techniques*. Technical bulletin no. 18. Ministry of Agriculture, Fisheries and Food, London, UK.
- Makkar HPS 2003. *Quantification of tannins in tree and shrub foliage*. A laboratory manual. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Makkar HPS, Becker K, Abel H and Szegletti C 1995. Degradation of condensed tannins by rumen microbes exposed to quebracho tannins (QT) in rumen simulation technique (RUSITEC) and effects of QT on fermentative processes in the RUSITEC. *Journal of the Science of Food and Agriculture* 69, 495–500.
- Merchant M and Riach DJ 1994. The intake and performance of Cashmere goats grazing sown swards. *Grass and Forage Science* 49, 429–437.
- Min BR, Hart SP, Miller D, Tomita GM, Loetz E and Sahl T 2005. The effect of grazing forage containing condensed tannins on gastro-intestinal parasite infection and milk composition in Angora does. *Veterinary Parasitology* 130, 105–113.
- Mueller-Harvey I 2006. Unravelling the conundrum of tannins in animal nutrition and health. *Journal of the Science of Food and Agriculture* 86, 2010–2037.
- Niezen JH, Charleston WAG, Hodgson J, Miller CM, Waghorn TS and Robertson HA 1998. Effect of plant species on the larvae of gastrointestinal nematodes which parasitise sheep. *International Journal for Parasitology* 28, 791–803.
- Osoro K, Benito-Peña A, Frutos P, García U, Ortega-Mora LM, Celaya R and Ferre I 2007a. The effect of heather supplementation on gastrointestinal nematode infections and performance in Cashmere and local Celtiberic goats on pasture. *Small Ruminant Research* 67, 184–191.
- Osoro K, Mateos-Sanz A, Frutos P, García U, Ortega-Mora LM, Ferreira LMM, Celaya R and Ferre I 2007b. Anthelmintic and nutritional effects of heather supplementation on Cashmere goats grazing perennial ryegrass–white clover pastures. *Journal of Animal Science* 85, 861–870.
- Paolini V, De La Farge F, Prevot F, Dorchies Ph and Hoste H 2005. Effects of the repeated distribution of sainfoin hay on the resistance and the resilience of goats naturally infected with gastrointestinal nematodes. *Veterinary Parasitology* 127, 277–283.
- Russel AJF 1990. Body condition scoring of goats. *Scottish Cashmere Producers Association Newsletter* 10, 3.
- SAS 1999. SAS OnlineDoc<sup>®</sup>, version 8. SAS Institute Inc., Cary, NC, USA.
- Silanikove N, Gilboa N and Nitsan Z 1997. Interactions among tannins, supplementation and polyethylene glycol in goats given oak leaves: effects on digestion and food intake. *Animal Science* 64, 479–483.
- Smith HA, Zoetendal E and Mackie RI 2005. Bacterial mechanisms to overcome inhibitory effects of dietary tannins. *Microbial Ecology* 50, 197–205.
- Stromberg BE 1997. Environmental factors influencing transmission. *Veterinary Parasitology* 72, 247–264.
- Tavendale MH, Meagher LP, Pacheco D, Walker N, Attwood GT and Sivakumaran S 2005. Methane production from *in vitro* rumen incubations with *Lotus pedunculatus* and *Medicago sativa*, and effects of extractable condensed tannin fractions on methanogenesis. *Animal Feed Science and Technology* 123–124, 403–419.
- Theodorou MK, Williams BA, Dhanoa MS, McAllan AB and France J 1994. A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. *Animal Feed Science and Technology* 48, 185–197.
- Van Houtert MFJ and Sykes AR 1996. Implications of nutrition for the ability of ruminants to withstand gastrointestinal nematode infections. *International Journal of Parasitology* 26, 1151–1167.
- Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3597.
- Van Wyk JA, Cabaret J and Michael LM 2004. Morphological identification of nematode larvae of small ruminants and cattle simplified. *Veterinary Parasitology* 119, 277–306.